D

in

in

as 18

ek

6,

ese

Public Health Reports

VOLUME 64

JUNE 24, 1949

NUMBER 25

IN THIS ISSUE

Rabies Problems and Control Pilot Mental Health Clinic



FEDERAL SECURITY AGENCY

PUBLIC HEALTH SERVICE

FEDERAL SECURITY AGENCY Oscar R. Ewing, Administrator

PUBLIC HEALTH SERVICE Leonard A. Scheele, Surgeon General

Division of Public Health Methods G. St. J. Perrott, Chief of Division

CONTENTS

	Page.
Rabies problems and control. A nation-wide program. James H. Steele and Ernest S. Tierkel.	785
Pilot mental health clinic. First annual report of Prince Georges County clinic. Mabel Ross	797
Notifiable diseases, year 1948	802
INCIDENCE OF DISEASE	
United States:	
Reports from States for week ended June 4, 1949 Territories and possessions:	807
Puerto Rico-Notifiable diseases-4 weeks ended May 28, 1949.	810
Plague infection in New Mexico and Utah	810
Deaths during week ended May 28, 1949	810
Foreign reports:	
Canada—Provinces—Notifiable diseases—Week ended May 14, 1949—World distribution of cholera, plague, smallpox, typhus fever, and vellow fever—	811
Cholera	811
Plague	812
Smallpox	813
Typhus fever	814
Yellow fever	815

Public Health Reports

Vol. 64 • June 24, 1949 • No. 25

Rabies Problems and Control

A Nation-wide Program

By JAMES H. Steele, D.V.M., M.P.H., and ERNEST S. TIERKEL, V.M.D., M.P.H.*

Few events arouse more terror in a community than the cry of "mad dog." The fear and anxiety which accompanies this disease has hovered over the country since the days of colonization. Records indicate that rabies was present in the colonies as early as 1753, and by 1860 the disease had found its way into most of the States east of the Mississippi River. It then moved westward with the growth of the Nation, and finally invaded California in 1899 (1).

ge.

85

02

07

10

10

10

11

11

12

13

14

15

In the last 15 years there has been an alarming increase in the incidence of rabies throughout many parts of the country. Its effects on health, agricultural economy, and wild-life conservation are disastrous. Each year over 30,000 persons are required to take the long and often painful series of vaccine inoculations as a result of exposure to rabid or suspected dogs. The annual cost to the country for human vaccine treatments and livestock losses exceeds \$5 million.

Surveys of present rabies control activities throughout the Nation on State and local levels have indicated a lack of uniformity which has minimized the effectiveness of individual control activities (2). Many communities have demonstrated effective rabies control programs; however, neighboring communities may have ineffective programs, or none at all. Epidemics may thus be reintroduced into rabies-free areas. Where one State employs one type of control program, the State to the north of it may employ another which conflicts with the strategy and tactics of its neighbor, while the State to the west may have no planned program at all. The hard fact is that the rabid animal respects no border lines, State or county, but is driven by pathological impulse to roam for miles, causing the spread of an epidemic from one area into another.

The Subcommittee on Rabies, National Research Council in 1945 (1) and the National Conference on Rabies in 1947 (3) unanimously

^{*}From the Veterinary Public Health Division, Communicable Disease Center, Public Health Service,

agreed that nation-wide uniformity of control procedures will be necessary for the eradication of rabies from the United States, and that this can be achieved only if a properly authorized national agency assumes the responsibility of coordinating rabies control activities.

A proposal has been made to form a Federal rabies control advisory committee composed of members from the Public Health Service, Bureau of Animal Industry, and Fish and Wildlife Service. The principal objective of this inter-departmental policy committee will be to draw a pattern of uniform control methods, based on the latest scientific information, for adoption and action by the States. In the operation of this national policy, the Public Health Service has agreed to (1) distribute to the States information on the latest accepted diagnostic techniques; (2) institute an accurate system of reporting; (3) keep local control authorities posted on the most effective immunization techniques; (4) assist in the drafting of licensing and dog-control ordinances, and (5) prepare and distribute educational material to insure wholehearted cooperation by the general public.

The Public Health Service has embarked on a functional plan for participating in the national program by offering its technical and administrative services to the States for a coordinated attack on the

rabies problem.

Serving as the nerve center for rabies control activities is the Rabies Control Branch, Veterinary Public Health Division, Communicable Disease Center. This Branch, with headquarters at Montgomery, Alabama, is active in investigations directed toward all aspects of the control of rabies. These investigations include the improvement and standardization of laboratory diagnostic techniques; training of State and local public health laboratory personnel through organized practical short courses; study of the immunology and pathology of the disease; testing and improvement of new experimental vaccines for animal immunization; preparation and distribution of educational material; epidemiological evaluation of reservoirs of infection; operation of field demonstration control projects; the furnishing of aid in epidemics, and the provision for consultation services in the development of permanent and long-range rabies control programs at the State and local levels.

Laboratory Diagnosis

The attack against any mass disease problem must necessarily begin with adequate diagnostic services. At the Rabies Control Branch laboratory studies are in progress on the comparative evaluation of various diagnostic techniques. The objectives are to determine the most efficient methods of making a diagnosis with the utmost accuracy, speed, economy and practicability. For example, in the

mic sur tole pre sim

me

of be cate place slide sprewhite slide

pres imp cent stru

Neg

of s com purp Selle stain limin prob

Si dyin strat surv perc miss It

furni tests giver of hu rate

Th the w

at ev

microscopic examination of animal brain tissue for Negri bodies, trial surveys have shown that the cumbersome practice of preparing histologic sections is not necessary. A film of brain tissue when properly prepared is just as accurate for purposes of diagnosis and is manifestly simpler, quicker and less costly.

y

В,

e

e

t

d

ol

0

r

d

e

8

e

d

ıl

e

The laboratory has also found that the three most acceptable methods for the application of the suspected brain tissue on the microscopic slide are: first, the so-called rolling technique in which a piece of brain tissue is rolled over the surface of the slide with a wood applicator; second, the smear technique in which a small piece of tissue is placed on one end of the slide, and is crushed with the aid of another slide and drawn down the length of the slide, creating a homogeneous spread of tissue; third, and perhaps best, the impression method in which a small section of tissue is placed on blotting paper and the slide is applied to the cut surface of the section with just enough pressure to leave a thin film of tissue imposed on the slide. In the impression technique an optimum amount of nerve tissue can be concentrated in a small area without damage to neuronal and interstitial structure (4, 8, 9, 10, 11).

A multitude of differential stains for use on brain-tissue smears for Negri-body examination are being used routinely with varying degrees of success in diagnostic laboratories throughout the country. In comparative studies on most of the staining techniques used for this purpose, we have achieved by far the most satisfactory results with Sellers' stain. Because the methylene blue and basic fuchsin of this stain are dissolved in methyl alcohol, the tissue film requires no preliminary fixation; it is fixed and stained simultaneously. This is probably the most important factor in making the Sellers' technique the most rapid and the most easily handled (δ) .

Since Negri bodies cannot always be found in the brains of animals dying of rabies, it is important that animal inoculation for demonstration of the virus be done on Negri-negative specimens. Extensive surveys of large numbers of rabies cases have shown that 10 to 12 percent of those cases proved positive by mouse inoculation had been missed by direct smear microscopic examination for Negri bodies.

It is therefore strongly recommended that laboratories which furnish rabies diagnostic services be equipped to do animal inoculation tests on Negri-negative brain tissues. In this way, vital assistance is given the physician handling a dogbite case where there is a question of human exposure, and necessary support in the form of more accurate reporting is given to the rabies control authorities who are aiming at eventual eradication of the disease.

The operation is simple and inexpensive. The preferred animal is the white mouse since it is uniformly susceptible, low in cost and easily handled. The intracerebral inoculation of a suspension of infected brain material will produce typical and constant symptoms in 5 to 11 days with the consistent production of Negri bodies (8, 10, 12). The details of the test have been standardized for practical use at our laboratory and are available for distribution to all agencies responsible for the diagnosis of rabies (13).

do

col

an

cal

An

An

Sai

Ser

1

vac

scie

He

wel

is n

vac

thu

Roc

. N

mai

dem

sent

Rah

Mer

alar

By

mor

for t

men

treat

in to

prog

dog (

It w. Worl

Heal progr

and a of 70

perio

vacci

23,00

Adde

It

Animal brains shipped to diagnostic laboratories are often grossly decomposed on arrival, making it impossible to inject mice without the danger of introducing complicating bacterial infection. Suitable agents for treating suspensions of such contaminated material are 10 percent ether, 0.5 percent phenol, 1:5,000 merthiolate, pure glycerol or 500 units of sodium penicillin G per milliliter of tissue emulsion. All of these agents will kill the contaminating bacterial without affecting the virulence of the virus. Comparative evaluations of these agents in our laboratory have earmarked the penicillin treatment of brain suspensions as the preferred technique. Such suspensions are ready for inoculation within 30 minutes as compared to 2, 6, 6, and 48 hours for tissue emulsions treated with ether, phenol, merthiolate and glycerol, respectively.

Implementing the diagnostic reference services of the Rabies Control Branch, an organized short course in the laboratory diagnosis of rabies is offered twice each year at the laboratories of the Communicable Disease Center, Atlanta, Georgia. This training is open to all grades of employed laboratory personnel including directors and senior staff members. Although first consideration is given to the laboratories of State and local public health departments and other official agencies responsible for the diagnosis of rabies, applicants from hospitals and similar institutions are considered when vacancies occur.

This one-week refresher course is designed essentially for practical laboratory training. The students carry out the various procedures to gain proficiency and to keep abreast of new and improved methods. It is supplemented by lectures and demonstrations. Phases of the course covered include orientation and background, preparation of stains, gross brain dissection, tissue-film preparation and staining, mouse inoculation and symptomatology, mouse brain smears and staining, microscopy (the Negri-body and differential diagnosis—other virus inclusion bodies), cost and materials, review and unknowns.

Canine Rabies Vaccination

One of the principal projects at the laboratory of the Rabies Control Branch is the study of the efficacy of canine rabies vaccination. Current studies in the laboratory and analytical observations of field trials have produced satisfactory evidence that annual vaccination of

dogs is an essential means of controlling rabies and should be encouraged to augment other measures such as licensing, quarantine and the elimination of strays.

ted

to 2).

at

eies

slv

ble

are

on.

out

of

eat-

us-

nol,

bies

osis

ompen

tors

1 to

and

pli-

hen

tical

ures

ods.

the

n of

ning.

and

sis-

wns.

ntrol

tion.

field

on of

The following organizations and official agencies have endorsed canine rabies vaccination: American Veterinary Medical Association, American Medical Association, American Public Health Association, American Animal Hospital Association, United States Livestock Sanitary Association, National Research Council, Public Health Service, and Bureau of Animal Industry (3).

Those States, counties and municipalities which put the annual vaccination requirement into effect several years ago did so on a scientific basis. Evidence, presented by Johnson of the International Health Division, Rockefeller Foundation, after carefully planned, well-controlled experiments and field studies, showed that immunity is maintained at a high level for 6 months and is effective 1 year after vaccination (6). Our experience in the Public Health Service has thus far corroborated the findings of the 9 years of research by the Rockefeller Foundation.

Mass immunization of dogs has been used with excellent results in many communities throughout the country. One of the most forceful demonstrations of the effectiveness of canine vaccination was presented just a year ago during an outbreak of rabies in Memphis, Tenn. Rabies incidence, which had been at a substantial endemic level in Memphis and in Shelby County for years, suddenly began to reach alarming epidemic proportions in the late winter and spring of 1948. By March, positive animal cases were being reported at the rate of more than one a day. The number of cases was twice that reported for the same period in 1947, and four times as many as in 1946. Over 150 persons had undergone the full series of anti-rabic vaccine treatments, which in the first 3½ months exceeded the total number of treatments for any previous entire year. At that time we were called in to assist in the planning and operation of an emergency control program.

It was noted that the disease continued to spread in spite of a strict dog quarantine, adequate stray dog control and a good licensing law. It was estimated that about 8,000 dogs were vaccinated annually. Working swiftly and efficiently, the Memphis and Shelby County Health Departments set up machinery for a voluntary dog-vaccination program by mobilizing the practicing veterinarians of the community and alerting dog owners through every conceivable medium. A series of 70 emergency dog-inoculation clinics were operated over a 6-day period at strategic locations throughout the city and county. The vaccination charge was \$1 per dog, regardless of dose. As a result, 23,000 dogs were inoculated during the 6-day emergency program. Added to those vaccinated by veterinarians in their routine practice,

preceding and following the campaign, it is estimated that 80 percent of the dog population was immunized.

The results of the Memphis program were phenomenal. In the ensuing months, the positive cases began to drop until the last case of animal rabies and the last human anti-rabic vaccine treatment were reported in July. Both city and county remained entirely free of rabies until March 10, 1949, when the first rabid animal since July 22, 1948, was picked up at the city limits. Seven months without a single case of rabies was a new and refreshing experience for Memphis.

a

ir

th

th

al

to

su

th

R

re

an

on

ce

int

the

sei

loc

vei

acc

nei

me

ma

for

ing

adv pol

I

Fiel

cou

exte

bros

I

Despite the thorough dog control activities, it was not possible to eliminate rabies from the area until mass immunization was added in the emergency program of 1948. The results of this episode stand as dramatic testimony to the importance of canine vaccination.

Great progress has been made in recent years in the improvement of canine rabies vaccine. The product now in use is a 20-percent phenol and heat-killed brain emulsion vaccine. Its manufacture has been standardized by Federal regulation, and all canine rabies vaccine on the market today is required by the Bureau of Animal Industry to pass the stringent Habel mouse potency test of the Public Health Service.

Investigations in animal rabies vaccination are being continued at the laboratory of Rabies Control Branch, and experimental activities along these lines have been expanded to include studies in the duration of immunity, dosage for maximal immunity response, efficacy of newly developed vaccines inactivated with a variety of chemical substances and by ultraviolet irradiation, and safety and relative antigenicity of experimental live virus vaccines such as the chick embryo vaccine.

Reporting

One of the three primary resolutions adopted by the National Rabies Conference held in Philadelphia in 1947 was "that rabies in animals should be made a reportable disease and the information be properly analyzed and distributed to all the States." Following this resolution, arrangements were made by the Public Health Service to print information concerning incidence of rabies in animals, as well as in man, in Public Health Reports which reaches all diseasecontrol agencies in the country. State health officers were asked to include rabies in animals in their weekly telegraphic reports to the This important statistical information is Public Health Service. compiled and distributed so that each State will have a week-by-week picture of the extent and movement of the infection in neighboring and other States. This service has proved invaluable during the past year in notifying authorities of new foci of infection so that control work could be started before the disease had time to spread.

The State Program

One of the most vital functions of the Veterinary Public Health Division is the assignment of qualified public health veterinarians to State health departments throughout the country. These Public Health Service veterinary officers are responsible for the organization and development of sound rabies control programs in the States to which they are assigned. They establish the extent and limits of the States' problem and set into motion the mechanism of control according to the results of their surveys.

Many State health departments employ full-time veterinarians of their own who cooperate with the Public Health Service. It is urged that this arrangement be effected in every State where rabies is prevalent, since it has been shown that the only States which are beginning to achieve a measure of success are those with programs under the supervision of a qualified public health veterinarian.

These men, whether Federally assigned or State employed, are the keystones in the fight against rabies. Through the offices of the Rabies Control Branch and Veterinary Public Health Division, they report the progress of their respective programs, learn of the problems and activities of other State programs, and obtain latest information on the technical and administrative aspects of effective control procedures. Through State professional societies, they stimulate the interest of the private practicing veterinarians of the State and enlist the active support of practitioners in local control programs. They serve as liaison officers between State diagnostic laboratories and local health units.

In administering the State-wide program, the health department veterinarian coordinates the efforts of local control by encouraging accurate reporting; alerting counties on the presence of rabies in neighboring areas; appointing local rabies inspectors; improving methods for the shipment of specimens to diagnostic laboratories; making canine vaccines available where needed; surveying facilities for collection and impoundment of stray dogs; preparing and distributing educational material throughout the State, and, by frequent visits, advising and consulting with local control authorities on current policies and methodology of control practices.

The Local Program

The actual legwork of control operations is done at the local level. Field demonstrations have proved that local programs work best on a county-wide basis or on a multicounty unit basis according to the extent of the local health jurisdiction.

It is suggested that all local rabies control programs include three broad measures. The first is impounding and destruction of all stray

836139 49 2

ent

the ase ent

free nce

out

his.

to ded

and

ent

ent has

cine

strv alth

ued ivi-

the

effi-

emitive

nick

onal s in

a be

this

e to

well

ase-

d to

the

n is

veek

ring

past

atrol

and ownerless dogs. This requires the operation of a local pound or humane shelter where stray dogs may be kept for a specified number of days and, if unclaimed at the end of that period, humanely destroyed. Strays should be collected by teams of dog wardens and assistants using trucks with proper enclosures. The second is annual anti-rabies vaccination of all dogs. The importance of canine vaccination in an efficient control program is now a firmly established fact and needs only a well-coordinated educational campaign to bring this fact to the public. The third is registration or licensing of all dogs. Licensing of all dogs in a community is an important adjunct of a successful control program. If properly enforced, it serves to defray the expenses for the over-all control program, assures a reasonably accurate dog census, rids the area of ownerless strays, and places the responsibility of dog control activities squarely on the dog owner.

h

h

8

u

i

The foregoing are essentials of a successful control program. If there is a local ordinance with or without the benefit of State enabling legislation, it should be administered by the local health department and enforced by city and county police officials. The ordinance should contain enforcing clauses. It should be well conceived and simply drafted and contain all important details with regard to its operation. The health officials should strive to launch an effective educational campaign in conjunction with any regulatory measures. This latter

step is paramount for the success of the program.

Some additional provisions which may be incorporated into the program are:

1. Except in heavily populated cities, vaccinated dogs, when properly tagged, may be allowed at large 30 days after vaccination. The vaccine is not fully effective until the end of that time.

2. In the face of an outbreak of rabies in the community, a strict quarantine should be placed on all dogs, requiring that they be confined at home or kept on a leash when out of doors.

3. Dogs under 6 months of age, which are particularly susceptible to rabies and not as readily immunized as adult dogs, should be confined.

- 4. Biting dogs and suspected rabid animals should be impounded and held under observation for 12 to 14 days so that proper medical disposition of dogbite and suspected human exposure cases may be effected. Dogs exposed to rabies should be destroyed or kept confined under observation for 6 months.
- 5. In case of rabies outbreaks in wild animals, such as foxes, adequate trapping programs should be instituted in cooperation with the State wildlife conservation authorities.

It should be pointed out again that adequate diagnostic facilities and human anti-rabies vaccine should be made available by the local health unit or branch laboratory of the State health department. All cases of rabies in man and animals should be reported to the local health officials.

d or

nber

nely

and

nual

cina-

fact

logs.

of a

fray

ably

the

If

oling

nent

ould

aply

ion.

onal

tter

the

per-

The

rict con-

ible

on-

ded

ical

be

on-

de-

the

ties

cal

Vaccination as a prerequisite to registration and combining the two as a single operation is a sound idea and will make the control program simpler and less cumbersome. The dog owner will appreciate the fact that he has only one trip to make each year when he can have his dog vaccinated and registered at the same time. He should be issued a single official uniform tag for the dog's collar and a single official uniform certificate of vaccination. The certificate can be made in triplicate, with the original for the dog owner, one copy for the health department's registration files and one copy for the veterinarian. A single fee should be charged which will be low enough to cover all classes of dog owners and high enough to defray expenses of vaccination services and the operation of stray dog control activities.

Another suggestion is that dog inoculation clinics be operated by veterinarians duly authorized by the health department. The clinics can be held at strategic points throughout the city or county over a 1- or 2-week period. It has been found that the effective control of rabies is enhanced by the desired immunization procedure of getting as many dogs as possible vaccinated in the shortest possible period of time. Temporary neighborhood clinics have the added advantage of making it more convenient for people to get their dogs vaccinated.

Education

No matter how soundly a local control program is developed and no matter how well an ordinance is drafted, it still must be supported by the people if it is to succeed. This can be achieved only by a well-planned educational program. In speaking of rabies, Denison has stated "there is no disease about which the public is more misinformed" (7). It is in the province of all of us as health workers to dispel superstitions, prejudices and misinformation. People will be happy to cooperate if they learn that canine vaccination is of proved value, that dog-registration ordinances are for their own protection in combating a deadly disease and not just another devious form of government revenue, that the conception of "dog days" during the hot summer months has no bearing on rabies, that the use of the "mad-stone" is a superstitious fantasy, and that dogs do not just "get rabies" spontaneously but as the result of exposure to another rabid animal.

In planning campaigns for many communities, we have successfully used such media as newspapers, radio, television, placards, sound trucks, mimeographed schedules of clinic stations, printed pamphlets, church announcements, and talks before civic and school groups in which audio-visual aids were utilized. In this connection, the Pro-

June 24, 1949 794

duction Division in cooperation with the Rabies Control Branch of the Communicable Disease Center is engaged in the production of audiovisual media for disseminating information on rabies-control methods. Two sound filmstrips on rabies were released during the past year and copies of each have been widely distributed throughout the country. They have proved to be valuable adjuncts in local control operations.

The first filmstrip, "The Fight Against Rabies", acquaints the audience with symptoms of disease in animals and man; refutes popular misconceptions such as the influence of climate and season on rabies prevalence; stresses the 100 percent fatality rate of the disease; presents the hazards of an uncontrolled epidemic; gives mode of transmission, and suggests means by which citizens can cooperate with control authorities in combating its introduction and spread through a community. Many of the points on control are illustrated by simple dramatization. Forceful material for mass immunization of dogs is also presented. The filmstrip provides excellent orientation for health department workers as well as being an effective educational device for the layman.

The second filmstrip, "The Laboratory Diagnosis of Rabies", is being used as a training aid for public-health laboratory workers in bringing to them visual demonstrations of the most efficient and practical techniques in the laboratory diagnosis of rabies.

Now in the script stage of production is a new motion-picture film

on rabies eradication which may be released later this year.

The Wildlife Problem

In many sections of the United States the rabies problem is complicated by the presence of the disease in wildlife. Wild animal rabies is now present in the States of the Appalachian Range from New York to Georgia, westward in the Southern tier to eastern Texas, and in the Midwestern belt that extends from western Pennsylvania through Iowa. Most of the cases reported are foxes, but civet cats, skunks and coyotes are also reported.

Without the problem of wildlife rabies, eradication of this disease from the United States would be a substantially easier task. Although we have a good vaccine for prophylactic measures in domestic animals, mass immunization tactics cannot be applied to the susceptible wildlife. In areas where wildlife rabies exists, coordinated programs between public health and wildlife conservation departments can be effectively developed.

Organized trapping programs sponsored by State game and fish commissions assisted by the Fish and Wildlife Service have produced good results in smothering serious outbreaks of rabies in wildlife. "Encirclement" and "thinning out" trapping techniques are suggested by wildlife conservation authorities and are based on the observation

of the

udio-

hods.

and

ntry.

ions.

audi-

oular

abies

pre-

rans-

con-

gh a

mple

gs is

vice

rs in

and

film

ipli-

bies

ork

d in

ugh

inks

ease

ugh

stic

sus-

ted

art-

fish

ced

life.

ted

ion

that rabies epizootics occur when there is an overpopulation of foxes. The disease in foxes, when present in epizootic proportions, will run its course in a year or more, the resulting mortality reducing the number of foxes until they are no longer a source of danger. However, besides the devastating effect of the disease in the foxes themselves, there is always the danger of some affected animals leaving the area and setting off a chain reaction of infection in a susceptible animal population at some distant place. There have been a few instances on record where infected wildlife served as the link in the transmission of rabies between isolated animal populations. The objective in trapping procedures is to reduce the fox population more quickly than the disease will and thereby shorten the period of infection and the danger of its spread to other animals and man.

Additional Control Services

In addition to the services available by the Communicable Disease Center for the control of rabies, activities of the National Institutes of Health, Foreign Quarantine Division, and Public Health Service Regional Offices are serving to strengthen the national control program.

Those aspects of rabies and its control which relate to the human disease are being studied in the Institute of Microbiology, National Institutes of Health. Here basic information is being sought through research of the disease in man. Another important activity at the Institute is the routine potency testing, improvement and establishment of minimum requirements for rabies biologics production.

At the end of the war, the Armed Forces requested that the Public Health Service include animal rabies provisions in their quarantine regulations. These regulations, now in effect, are administered by the Foreign Quarantine Division and require that all animal pets coming from countries where rabies is known to exist be vaccinated not more than 6 months prior to debarkation at the port of entry.

Throughout the country, Public Health Service Regional Offices are available for consultation on rabies and other disease control problems. Regional office staffs stimulate reporting, and encourage and assist in coordinating control activities in the regions. For States which require financial aid in the development of rabies control programs, grants-in-aid funds are available from the Public Health Service under P. L. 410, sec. 314c, 78th Congress for the operation of general health programs. These grants to the States are administered by the Regional Offices.

Conclusion

The eventual eradication of rabies from the United States is not an unobtainable goal. Its achievement can mark one of man's great victories over disease. The nature and extent of the problem is well known, the implements for attaining it are available and the strategy has been planned. It is felt that the integration of the services outlined in the foregoing pages with those of the State and local rabies control agencies in a unified national program will provide the modus operandi to reach this goal.

REFERENCES

- (1) National Research Council: Report of the Subcommittee on Rabies, Nov. 26,
- 1945. J. Am. Vet. Med. Assoc. 108: 293–302 (1946).
 (2) New York Academy of Medicine: Control of rabies. Report by the Com-
- mittee on Public Health Relations. Pub. Health Rep. 62: 1215 (1947).

 (3) Report of the National Rabies Conference, Philadelphia, 1947, Am. J. Pub. Health 38: 97–99 (1948).
- (4) Tierkel, Ernest S.: Inauguration of rabies control studies by the U. S. Public Health Service. J. Am. Vet. Assoc. 112: 18-24 (1948).
- (5) Sellers, T. F.: A new method for staining Negri bodies of rabies. Am. J.
- Pub. Health 17: 1080 (1927).

 (6) Johnson, H. N.: Experimental and field studies of canine rabies vaccination. In Proceedings of the Forty-ninth Annual Meeting of the U. S.

- tion. In Proceedings of the Forty-ninth Annual Meeting of the U. S. Livestock Sanitary Association, 1945, pp. 99-107.
 (7) Denison, G. A. and Dowling, J. D.: Rabies in Birmingham, Ala. J. A. M. A. 113: 390 (1939).
 (8) Webster, L. T. and Dawson, J. R.: Early diagnosis of rabies by mouse inoculation. Proc. Soc. Exper. Biol. & Med. 32: 570-573 (1935).
 (9) Johnson, H. N.: Significance of the Negri body in the diagnosis and epidemiology of rabies. Illinois Med. J. 81: 382 (1942).
 (10) Leach, C. N.: Comparative methods of diagnosis of rabies in animals. Am. J. Pub. Health 28: 162-166 (1928).
- J. Pub. Health 28: 162-166 (1938).

 (11) Damon, S. R. and Sellers, T. F.: A note on the probability of error in the diagnosis of rabies by microscopic search for Negri bodies. J. Lab. & Clin. Med. 27: 71-74 (1941).
- (12) Hoyt, A. and Jungeblut, C. W.: Experimental rabies in white mice and attempted chemotherapy. J. Infect. Dis. 74: 418 (1930).
 (13) U.S. Public Health Service: Mouse inoculation test for rabies. A laboratory
- diagnostic procedure for Negri negative and questionable brain specimens. (Communicable Disease Center, Atlanta, Ga., Mimeographed, 1948).

Pilot Mental Health Clinic

gy ut-

ies lus

26,

m-

ıb.

S.

J.

8.

1.

se

1-

1.

e

d

First Annual Report of Prince Georges County Clinic

By MABEL ROSS, M. D.*

The Prince Georges County Mental Health Clinic is a demonstration clinic under the National Mental Health Act and is financed by congressional appropriation for this purpose. It was established in January 1948, in Prince Georges County, Maryland, which includes suburban and rural areas in its 482 square miles and has a population of over 150,000.

The clinic was set up at the request of citizens and organizations of the county, who, with the county and State health departments, worked out the details of organization with the Mental Hygiene Division of the Federal Public Health Service. A group of interested citizens formed an interim board and, among other things, made the arrangements between the county and the University of Maryland to have the clinic housed in a building on the university campus. This group then organized a meeting of the representatives of organizations in the county which were interested in mental health. Under a constitution adopted at this meeting the representatives formed a committee-at-large to meet annually and elect the members of the advisory board. The 12-member advisory board of the Mental Health Clinic meets monthly with the clinic staff in an advisory capacity and acts as liaison with the community.

This pilot clinic was established to study methods through which the mental health program can be made part of the over-all health program for a county and to determine methods which communities can apply to attack the factors contributing to the high incidence of psychiatric disorders. Toward this end it is necessary to service individual cases in order to demonstrate the value of psychiatric treatment in already existing disorders and also to develop confidence in the mental health principles stressed by the clinic. Behind and beyond the individual service, each case serves as a basis for study and research of the community factors contributing to the existing illness and the possible points at which preventive action might have made psychiatric treatment unnecessary. As an important part of the treatment of those referred to the clinic, the existing community facilities are given an active role in the treatment plan. This activity has definite educational value and makes available the basic principles of mental health to a wider group than just those patients who are referred to the clinic.

^{*}Senior surgeon, Public Health Service, and Director, Prince Georges County Mental Health Clinic, College Park, Md.

798 June 24, 1949

The Prince Georges County Mental Health Clinic opened officially January 19, 1948, although one of the psychiatric social workers had been working with the County Health Department for the preceding 8 weeks. The present staff consists of two psychiatrists, two psychiatric social workers, one clinical psychologist, one mental health nurse, and three clerical staff members. The clinic sees patients only by appointment, which can be applied for by any resident of the county.

The activity of the clinic is divided into three parts for the purpose of this report: individual service and treatment; community activities;

demonstration.

Individual Service and Treatment

From January 19, 1948, through January 18, 1949, 423 patients were seen by the Prince Georges County Mental Health Clinic. these, 281 were children under 18 years, and 142 were adults. These patients were referred by physicians, schools, the probation department, social agencies, ministers, and other local groups as well as by friends and relatives. Some of them came on their own initiative. Twelve and one-half percent were referred by local physicians who frequently carried on treatment of these patients in collaboration with the mental health clinic; 37½ percent were referred by friends, relatives, or came on their own initiative. Many learned of the clinic from publicity in newspapers and magazines, but the majority learned of it from other patients. At the end of the year there was a waiting list of 31 patients to be seen as soon as appointments could be arranged. All of the time allotted for examination and treatment is now being utilized, and the rate at which patients are discharged is somewhat less than the rate at which applications are being made. Treatment time is concentrated on individuals whose problems are interfering with satisfactory functioning at work, home, or school. The seriously ill are referred to other facilities for the necessary intensive psychiatric treatment.

During the year there were 18 cases which would have required care in a general or mental hospital or in an institution had it not been for the clinic. Four patients who had attempted suicide are now making a reasonably good social adjustment. Five potential divorces have thus far been avoided and in at least 16 cases the economic status of the family has definitely improved either because the patients were able to accept employment or change to full-time employment. In almost all instances this was not due solely to treatment at the clinic but was accomplished with the cooperation of appropriate community

facilities.

As a result of the study of the cases referred to the clinic we are convinced that some of the more serious cases could have been prevented had there been available in the past adjustment clinics in the schools, better guidance of parents through pediatricians or public health nurses, and improved child welfare services to deal with early problems of foster home or other placement of children.

Community Activities

The relationship between the clinic and the health department of Prince Georges County existed before the actual opening of the clinic. To study ways of including the mental health program in the general health program of the county, the mental health nurse of the clinic has carried out several activities with the cooperation of the staffs of the mental health clinic and the health department. Liaison with the supervisory and staff nurses of the health department included conferences concerning patients to be referred to the mental health clinic or problems in their own case load. Participation in the health department in-service educational program has been through mental hygiene conferences and by the attendance of public health nurses at the mental health clinic staff conferences.

Other activities included organization of mothers' discussion groups in prenatal and well baby clinics in the county. These groups of 6 to 10 mothers have met weekly or monthly to discuss various aspects of child care and development. Individual conferences were held with mothers and fathers attending the well baby clinic, with discussions centering around questions brought by the parents, such as: "Why does my 2-year-old take things away from the 6-month-old baby?" "Will my baby be 'spoiled' if I pick her up when she cries?" "My little girl, age 3, will not eat all of her food. Does she need a tonic?"

Recently, conferences were held with the public health nurses, and they indicated an interest in extending such discussion groups as part of the general health program. This is in keeping with the belief of the health department and the clinic that neither the patients nor the staff should think of mental health as separate from general health. The aim of a mental health demonstration is to be so incorporated into all health, educational, industrial, and community thinking that it is no longer recognizable as an entity.

To study the problems of a traveling clinic in a strictly rural area, a part-time clinic was established in November 1948 in Upper Marlboro, where the transportation problems are greater. This clinic has been held regularly twice a month.

The schools have sought help in individual cases and also have brought group problems for consultation. Discussion between the clinical psychologist and school representatives concerning individual children referred to the clinic, led in the summer of 1948 to setting up

ly

id

ng

i-

e,

7.

3;

y

June 24, 1949 800

a reading class by the county department of education with the cooperation of the mental health clinic. As a result, the schools have set up some classes and plan to continue the summer sessions. Other problems of maladjusted children needing special education are being studied by the clinic and schools. Conferences were held in five schools, and the referred problems were used as the starting points for discussion. These were enthusiastically received.

The Parent-Teachers Association groups requested assistance in county institutes. They feel it is their responsibility to make teachers and parents aware that help is available for some of the school

problems.

Although the courts have referred few cases, it has been possible to set up a good working relationship with the probation department. Despite being seriously overloaded, the men of this department have taken time to discuss cases before and after referring them to the clinic.

Other community activities included consultative conferences with the Welfare Board, Catholic Charities, Social Service League, Vocational Rehabilitation Service, and the Maryland State Employment Service.

The clinic staff has made many speeches to service clubs, Parent-Teachers Association, women's groups, university classes, home-makers' representatives, the Community Council, and the Medical Society. These talks almost invariably were followed by periods of discussion. Recognizing that all interested organizations in the county now have a working knowledge of the clinics, it is felt that they want more than a speech. For this reason, the clinic staff will try to work with the interested group in panel discussions, institutes, and working conferences planned toward action more than discussion of the clinic itself.

Demonstration

During the past year the clinic has had professional visitors from other States and other countries as well as from its own county and other parts of Maryland. Among them were psychiatrists, psychologists, nurses, social workers, and health officers. They came from private hospitals, other county clinics, State hospitals, and administrative departments. They came from New York, Virginia, West Virginia, North Carolina, Illinois, Iowa, Colorado, Texas, Arizona, and Oregon. They also came from China, Colombia, and Scotland.

While the visitors were interested in the functioning of the clinic per se, practically all hoped the demonstration clinic would be able to answer the problems in their own situations. Most of their questions centered around the community relations of the clinic—what agencies are worked with directly, both on a referral and a

the

ave

her

ing

nts

in

ers

ool

ble

nt. ent

to

ith

ca-

ent

at-

10-

cal

of

he

at

ill

es.

on

m

nd

0-

m

8-

st

a, d. ic le ir

a

consultative basis; what were the extramural activities, both consultative and educational; what were the mechanisms of cooperation with the community agencies; how can cooperation with the local medical society be established and maintained; how can the interest and support of the community and local agencies be fostered?

There have been nonprofessional visitors also. Telephone inquiries regarding the clinic come almost daily—usually from out of the county—and here an attempt is made by the receptionist to direct the caller to other clinics. Publicity has also brought letters from all over the country—some curious and some appealing. Each letter has been answered, usually with referral to the State Mental Health Authority for information as to nearby clinics. In two instances patients suddenly appeared—one from Pennsylvania (referred to and aided by a mental hygiene clinic in that State) and one from Ohio (referred to the State Mental Health Authority). Most of the publicity resulted from an article in a national women's magazine stressing the community participation in planning the clinic and the individual service.

There also were letters from cities and counties in North Dakota, South Dakota, Ohio, Wisconsin and California, requesting information about setting up similar mental health clinics.

It appears that this clinic has begun to serve its function as a demonstration to other counties in other States throughout the coun-With every bit of progress, wider possibilities open for com-With every report there is increasing evidence of the munity activity. great numbers of people throughout the country who are seeking some pattern and direction in establishing mental health services. Many questions have been about community cooperation, and it has been a source of pride to the clinic staff to be able to stress the active part played by the people of Prince Georges County in having the mental health clinic set up in this county, and to point to the continued interest and active support of the advisory board. However, the most unusual feature of this demonstration clinic is the committeeat-large, representing all interested groups in the area and expressing a community sense of responsibility. We hope that, with the continued help of the people in the county, more community methods will be found to serve as patterns for the development of other community-wide programs for mental health.

Notifiable Diseases, Year 1948

by law or regulation to be reported and the figures are included although manifestly incomplete. There are also variations among the States in the degree of, and checks on, the completeness of reporting of cases of the notifiable diseases; therefore comparisons as between States may not be justified for certain diseases. As compared with the deaths, incomplete case reports are obvious for such diseases as The figures in the following table are the totals of the monthly morbidity reports received from State health authorities for the year These reports are preliminary and the figures are more or less incomplete and subject to correction by final reports. The figures The comparisons made are with similar preliminary reports; but, owing to population shifts in many States since the 1940 census, the figures for some States may not be comparable with those for prior years, especially for certain diseases. Each State health officer has been requested to include in the monthly report for his State all diseases that are required by law or regulation to be reported in the State, Only 11 of the common commalaria, pellagra, pneumonia, and tuberculosis, while in many States other diseases, such as puerperal septicemia, rheumatic fever, and may be assumed to represent the civilian population only, although in some instances a few cases in the military population may be included. municable diseases are notifiable in all the States. In some instances cases are reported, in some States, of diseases that are not required although some do not do so. The list of diseases required to be reported is not the same for each State.

annually in consolidated form, have proved of value in presenting early information regarding the reported incidence of a large group of diseases and in indicating trends by providing a comparison with similar preliminary figures for prior years. The table gives a general In spite of these and other deficiencies inherent in morbidity reporting, these monthly reports, which are published quarterly and picture of the geographic distribution of certain diseases, as the States are arranged by geographic areas. Vincent's infection, are not reportable.

Leaders are used in the table to indicate that no case of the disease was reported.

Consolidated monthly State morbidity reports for the year 1948

	Pneu- monla, all forms	706 72 70 70 1 264 1,858	3, 180	2,445 1 2,445 2,776 412 412
	Pella			
	Oph- thal- mia	77.7	⊉ ∞∞	471
	Mumps	962 432 579 14, 216 4, 441	9, 603 32, 152 15, 598	3, 363 11, 704 13, 202 18, 681
	Men- ingitis, menin- gococ- cal*	- 510 E12	22.92	136 220 106 78
94	Mea-	2,967 1,228 1,648 37,108 623 3,281	57, 776 42, 855 39, 185	26, 755 41, 796 37, 273 37, 273
r 1940	Ma- laria 1	23 10 17	36	-5881
the year	Influ- enza	40 40 13 86 86	- 233 207 128	121 408 148 78 1, 215
Jor	Hook- worm disease	1	• 113	4.88
reports	Ger- man mea- sles	153 46 1,076 23 356 356	2, 128	228 228 545 1,170
pranty	En- cepha- litis, infec- tious	8 - 2 - 0	36.31	2000
te mor	Dysen- tery, unde- fined			10 8
ty State	Dysen- tery, bacil- lary	122 10 10 7	371	42 19 146 117
month	Dysen- tery, amebic	402	200 E1	74 14 338 518 14
dated	Diph- theria•	303 303 20	328	340 433 127 127 40
nosuo	Con- juncti- vitis i	178	48	10 3 151 171
	Chick- enpox	3, 374 790 2, 983 18, 127 1, 555 10, 054	26, 523 27, 098 26, 820	18,004 3,638 15,925 20,478 33,513
	An- thrax	010801	168	00000
	Division and State	Maine Maine New Hampshire New Hampshire Massachusetts Rhode Island Connectiout.	New York New Jersey Pennsylvania. BAST NORTH CENTRAL.	Ohio

######################################	2 2 3 2 4 4 6 5 1 1 2 3 2 4 4 6 1 1 2 3 2 4 4 6 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1, 5, 2, 4, 5, 1,	1, 652 1, 863 1,058 10, 863	263 322 1,060 1,615 201 40	1,046	80, 072 77, 995 777, 901	16.28
	. 4 9 2801	12.13	193	11 2	0 0 0 0 0 0 1 0 0 1 0 0 1 0 0 1 0 0 1 0 0	1,001	
	8 2 2 2	8 52	8 10 10 10	-1 001-		1,078	
3,115 5,550 144 729 3,100	4, 967 230 1, 967 1, 048 1, 329 1, 329	1,306 1,132 818 371	808 225 889 7, 815	2,2,808 2,720 6,57 1,282 1,963 1,963	12, 246 2, 595 32, 483	238, 764 155, 852 175, 643	3, 251
8885288	*******	5858	22.002	811780440	328	3, 375 3, 399 8, 035	P-M
9,330 10,151 6,176 1,518 1,014 3,834 1,354	12, 334 73, 746 7, 746 7, 746 7, 173 7, 173 4, 802	4, 410 2, 042 1, 304	4, 327 1, 542 1, 626 51, 121	10,946 1,222 1,222 1,222 2,533 384	12, 914 6, 436 65, 069	613, 810 221, 115 612, 068	2, 420
828 -64	39 30 3,668 111	80 319 120	83, 403 8, 594	-	2000	9, 797 17, 317 53, 575	1
2222 255	73 2, 200 2, 200 1, 296 1, 296 1, 366	2, 363 6, 924 745	7, 795 1, 137 3, 971 58, 604	475 983 33 1, 604 11, 225 11, 225 174	3, 267 2, 940 14, 153	165, 805 443, 339 452, 101	300
10	1,917 5,168 5,008	3,210	242	6		16, 260 12, 972 15, 236	6
7 11	236 236	88 38	10 ET	107 187 188 188 188 188 188 188	4,280	18, 287 12, 667 38, 339	22.12.33
204488 E	84 846	හනුගන	-10000	N +889 8	9002	878 889 880	8 0 9 0 9 0 9 8
2 88 4	143	75	1, 237	5 5	164	15,085 9,516 9,421	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
110	82 01 82 171 172 173 173 173 173 173 173 173 173 173 173	87 106 192	20,081	305	35	23, 727 16, 979 30, 872	988
12 89 12 4	22.23	314 36 100	361.	-10800000	23 149 315	4, 510 3, 130 3, 341	0.0
¥88888	180 203 203 503 606 606 827	357 292 614 266	1362	27 27 27 28 28 28 28 28 28 28 28 28 28 28 28 28	91 75 443	9,610 12,405 14,943	63-63
31	88.84	71	0 8 E E 0 1 E E 0 2 E E 0 2 E 0 0 E E 0 0 E E 0 0 E E 0	28 48 2	218	1, 458 1, 863 1, 329	21
3, 4639 3, 460 3, 264 1, 776 5, 075	3, 686 3, 686 1, 196 1, 629 1, 402	1, 190 2, 080 1, 430 1, 430	1,891 354 1,418 14,563	2, 557 1, 242 2, 200 3, 889 191	8, 620 1, 862 40, 539	331, 431 317, 565 301, 423	1.941
0000-00	-0000000	-000	0-00	000000	101	4778	0 A A A A A A A A A A A A A A A A A A A
Minnesota. Jowa. Missouri. North Dakota. South Dakota. Kansas. SOUTH ATANTIC	Delaware Maryland District of Columbia Virgina West Virgina North Carolina South Carolina Georgia Florida RAST SOUTH CENTRAL	Kentucky Tennessee Alabana Mississippi WEST SOUTH CENTEAL	Arkansas Louisians. Oklahoma Tenas.	Montana Glasho W yoming Colorado Colorado Arizona Utah New Maxico	Washington Oregon California	Year 1947 Median 1943-47	Alaska Hawali Territory Panama Canal Zone is

5 1,170 1,215 11 37,273 78 18,681 5 5412

40

0 33, 513

Wisconsin

See footnotes on pages 805 and 806

Consolidated monthly State morbidity reports for year 1948—Continued

Whoop- ing Cough•	1, 326 2, 772 330 724	4,1994	2, 29, 98, 98, 98, 98, 98, 98, 98, 98, 98, 9	E 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	288 94 1.9. 288 289 1.9. 288 288 288 288 288 288 288 288 288 288
Vin- cent's infec- tion	284 3		128	35 88 88	14 6
Unda- laut fever*	. 2032-10	25 88	PE 212 200	22 4 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	444845 8 55
Ty- phus fever, en- demic	N	10	-	9 1	228 228 228 170
Para- ty- phoid fever	25.001	1 20 33	17 7 10 1139 14 139	20-22-4	4224558
Ty- phoid fever*	1122 % 252	186 38	28.88.25.	188888	**************************************
Tula- remia	100	-0	44404	844 888	e∞3444888
Tuber- culosis, respir- atory	478 2, 814 483 1, 486	118,777	2, 346 6, 447	325	2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2
Tuber- culosis, all forms*	2, 996 1, 908 1, 908	14, 623. 3, 136 4, 940	4, 717 6, 950 6, 109 8, 109	3,307 2,067 346 231 471 1,108	8, 8, 8, 8, 8, 8, 8, 8, 8, 8, 8, 8, 8, 8
Trich- inosis	7 14 88	288	111 7	122	
Tra-	60	1	40 4	5 28 28	89 89
Teta- nus		828	248	1 1 9	r 84 03E
Small- pox*		1	0.00		
Septic sore throat	122812	E	30 37 77 515 195	18 12 12 13 13 14 15 15 15 15 15 15 15 15 15 15 15 15 15	-28 88 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
Scarlet fever*	630 167 200 6,053 346 1,094	2, 505 7, 795	10, 153 2, 024 4, 514 5, 934 2, 216	1, 629 1, 280 1, 125 137 1, 024	200 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
Rocky Moun- tain spotted fever	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	812	947		200 200 200 200 200 200 200 200 200 200
Rheu- matic fever	182	1,108	139 151 151 588	106	10 L1 81 82 10 82 10 10 10 10 10 10 10 10 10 10 10 10 10
Rabies in man		6 6 8 6 8 8 8 8 8 8 8 8 6 8 8 6 8 8	9 -		
Polio- myeli- tis*	11885 11885	1,402	1,173 397 1,101 651	1, 426 1, 200 1310 127 127 128 324	128 1156 173 2 518 2 518 2 518 2 518
Division and State	NEW ENGLAND Maine. New Hampshire. Vermont. Nassechusetts. Rhode faland. Ognnecticut.	New York New Jersey Pennsylvania. East North Central	Ohlo. Indiana. Illinois Michigan. W.Jaconsin.	Minnesota. Missouri North Dakota. North Dakota. Nobraska. Kansas.	Delaware Maryland District of Columbia Virginia Wet Virginia North Carolina South Carolina Georgia

3, 170 3, 106

Georgia

	1, 273 1, 255 1, 255 143	1, 267 250 1, 054 12, 053	346 289 1, 838 1, 236 27 27	1, 012 3, 710	74, 204 155, 991 132, 814	146 524 111
2000	88	п	~878 S	462	1, 618 2, 332 2, 332	9
	5282	2552	*818°58°	153	4, 886 6, 147 4, 959	1
	2777	34-25°		10	1, 184 1, 901 4, 517	21 17
	∞ ⊙ +≈	4128	10 10 10 10 10 10 10 10 10 10 10 10 10 1	118 118	1,0057	60-4
	88238	312	. 38887a	288	2, 905 3, 062 4, 221	800
	-828	2852	311	00 mt	1,093 1,360 887	8 2 6 6 1 6 5 3 6 6 3 6 6 3 6 6 3 6 6 6 6
	2,032	2, 393	2,280 1,480 1,480 1,480 1,480	8, 266	4 73,266 70,056 67,824	1,024
1	29.746 27.746 800 800	2, 427 3, 157 2, 230 13, 296	718 243 60 1, 561 19, 357 19, 255 19, 266	2, 360 8, 830 8, 886	133, 500 130, 474 118, 307	1,072 1,072 19 41
		10	•	8 8	461 428 317	1 1 1
Ī	2- 6	201 222	478 H2	17 88	2, 202 1, 103 1, 618	8
-	44å0	280		5	58.55 59.05	111
Denne C	4 5		- PD-04	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	+ 56 173 356	(D)
	\$8	1, 190 46 106 3, 598	88034488	103 216 580	19, 277 15, 906 9, 525	88-
	1,366 1,598 651 269	262 192 506 1, 296	510 218 1118 757 253 170 512 512	1,856 787 3,719	• 78,662 84,370 142,274	38-
	17 36 16	- 1900	81-38- 4-	10	526 565 452	6 6 E 6 8 E 8 8 6 6 9 6 6 9 6 6 8 8
	35 8	25.7	-2488521	88 797 897	4, 386	16
	*******			0 0 1 0 0 1 0 0 1 0 1 1 0 1 1 0 1 1 0 1 1	888	6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6
	* 202 201 162 162	145 156 362 1, 765	152225528	385 219 4 5, 857	4 27,902 10, 734 13, 514	1011
EAST SOUTH CENTRAL	Kentucky Tennessee Alabama Mississippi Wast south central	Arkansas Louisiana Oklahoma Texas	MOUNTAIN Montana Idaho Wyoming Colorado New Mexico New Mexico Arlrona Utah Nevada	Washington Oregon California	Year 1947 Median 1943-47	Alaska Hawaii Territory Panama Canal Zone ii.

Footnotes to Tables on Pages 802 to 805

Diseases marked with an asterisk..() are reportable by law or regulation in all the States; brinding the District of Columbia. Typhoid fever is reportable in all the States; paratyphoid fever in all accept 6 States. Syphilia is reportable in all the States and the District of Columbia but is not included in the table. Some States have increased and some have reduced the list of reportable diseases since the latest published compilation of reportable diseases since the latest published compilation of reportable diseases (Full. Reprint No. 2544.
Includes cases (Full. Reath Rep. 59: 317-340 (1944). Reprint No. 2544.
In a lew States practically all cases contracted outside the United States.

3 All cases reported as 'ophthalmia neomatorum since July 1, 1948. Prior reports included abuse croports

I Lobar pneumonia only.

Ne New York City only.

Exclusive of cases artificially induced.

Reports of chickenpox not received from Iowa for the months of July, August, September, and October, 1948 (inclusive) only.

Reported January-June 1948 (inclusive) only.

Reported January-June 1948 (inclusive) only.

Includes non-selfest case.

Contracted out-of-State.

In the Canal Zone only.

Footnotes to Tables-Continued

16 Includes septic sore throat. 16 Included in scarlet fever.

¹⁷ The report of a case of smallpox in the Panama Canal Zone for the first quarter of 1948 (Pub. Health Rep. 63; 98; (1948)) was an error. Also I case of typhus fever should have been recorded for the Canal Zone for that quarter.

The following list includes certain rare conditions, diseases of restricted geographical distribution, and those reportable in or reported by only a few States; last year's figures

in parentheses (where no figures are given, no cases were reported last year, or the disease was not included in last year's published tabulation.

Was not included in last year's published tabulation.

Minnesota & Go, South Dakoto 5, Nebraska i (1), Kentucky I, Tennessee I, Idaho 2, Nerada 3 (1), Hawaii Territory I (1), New York & Include 5 (1), New Mexico 9 (6), Oregon 3 (2), California 6 (6), Alaska 2

Cancer: North Dakota 790, Kanasa 3,882, South Carolina 1,460, Georgia 222, Florida Cancer: North Dakota 790, Kanasa 3,882, South Carolina 1,460, Georgia 222, Florida 1,860, Kentucky 22, Tennessee 2,917, Alabama 3,476, Missiskapi 1,485, Arkansas 677, Louisisna 2,130, Monthan 69, Idaho 831, New Mexico 643, Utah 271, Nevada 7, Coccidiolomycosis: Kanasa 1, Oklahoma 1, New Mexico 1, Arizona 15 (2), California

Colorado tiek fever: Colorado 67 (69). Dengue: Virginia 1, South Carolina 7 (14), Florida 1, Mississippi 1 (1), Oklahoma 1,

Teras 14 (19).
Dermatitis: New Hampshire 48 (15), Missouri 37 (117), Kentucky 150 (mycotic derma-

Diarrhea: Rhode Island 5, Connecticut 25, New York 170 (278), New Jersey 3 (41), Pennsylvania 28 (138) Includes gastroenteritis, Olio 1,311 (709) Includes enteritis, Indiana 20 (4) Includes enteritis, Illinois 54 (92), Michigan 101 (17), Iowa 1, Kanasa 48 (116) Includes enteritis, Maryland 36 (93), West Virginia 8 includes enteritis, South Carolina 11,856 (12,246), Florida 191 (68), Kentucky 17 (3) Includes enteritis, South Carolina 11,856 (12,246), Florida 191 (68), Kentucky 17 (3) Includes enteritis, Oklahoma 7 (2), Idaho 177 (2) includes enteritis, Colorado 24 (3) enteritis, New Mexico 54 includes enteritis, New Mexico 55 (694), Arkanasa 698 (594) includes all animals.

Sac 26,034), Arkanasa 698 (594) includes all animals.

Enceptalistis, other forms: Maine 1, Rhode Island 2, Connecticut 2, New York 7, Ohio 22, Michigan 67, Nebraska 1, Delsawae 1, Maryland 15, Florida 24, Kentucky 11, Adalor 7, Colorado 13, New Mexico 11, Arizona 1, Washington 12, Fanama Canal

Exystpelae: New Hampshire 4, Vermont 1, Connecticut 17, Ohio 36, Indiana 12, Illinois 122, Michigan 22, Wisconsin 60, Minacota 3, North Dakota 11, South Dakota 12, Louisiana 9, Oklahoma 4, Montana 10, Idaho 20, Colorado 50, New Mexico 3, Usha 2, Norada 1, Washington 16, Oregon, 30 Anaka 2, Hawaii Territory 12. Favus: Florida 1, Kantucky 6 (3). Ornecticut 1, New York 566, New Jersey 116 (20) includes cases reported as food infection, Ohio 13 (22), Indiana 6 (18), Illinois 89 (79) includes cases reported as food infection, Ohio 13 (22), Indiana 6 (18), Illinois 89 (79) includes cases reported as food infection, Minsota 30 (1144), Kansas 6, West Virginia 2, Florida 13, Louisiana 8 (30), Oklahoma 60 (8), Idaho 13 (6), Colorado 307 (7), New Mexico 27 (29), Washington 73 (481), Oregon 17 (48), California 67 (10, New Mexico 27), Missisipp 110 (306), Louisiana 174 (209), Idaho 3, Arisona 1 (1), Provincesee 60 (74), Missisipp 110 (306), Louisiana 174 (209), Idaho 3, Arisona 1 (1),

9, Tennessee 60 (74), Mississippi 19 Utah 1, Nevada 1, California 5 (7).

Impetigo contagiosa: Vermont 1, Rhode Island 3, New York 91 (71), Ohio 139 (32), Indians 53 (89), Illinois 53 (80), Michigan 1,425 (1,450), Missouri 6, 165), North Dakota 78 (6), Nobreaks 9 (4), Kansas 53 (58), Maryland 4 (10), Kentucky 39 (24), Montana 46 (40), Idaho 69 (80), Wyoming 25 (27), Colorado 77 (60), Newada 13 (152), Washington 87 (1,044), Alaska 11 (7), Hawaii Territory 96 (61).

Laundice (including hepstitis and Well's disease): Maine 11 (32), New Hampshire 2 (4), Rhode Island 1 (1), Connectiont 3, New York 211 (59), Pemasylvania 64 (49), Ohio 2 (7), Indiana 1 (6), Illinois 16 (27), Michigan 17 (13), Minnesota 42 (29), North Dakota 1 (12), Maryland 8 (8), South Carolina 2 (8), Florida 66 (29), Kentucky 72 (6), Washington 8 (17), Oregon 20 (79), California 107 (166), Hawaii Territory 20 (7), Fanama Qanai Zone 29.

Kala-azar: Montann 1, reported in April 1948. Contracted outside the United States. Leprosy. New York 9 (5), Florida 11 (2), Louisiana 7 (7), Tenza 10 (16), Arizona 1, California 15 (13), Hawaii Territory 27 (29), Panama Canal Zone 3 (1).

Lymphocytic edoriomeningtis: Maine 1, Massachusetta 19 (6), Khode Island 5, Indiana 1, Minnaeota 1 (6), Tennessee 11 (13) includes ohoriomeningtis undefined.

Lymphogranuloma venerum: New Hampshire 1, Connecticut 2, Missouri 8 (26), Florida 197 (216), Kentucky 4, Tennessee 68 (99), Mississippi 40, Louisiana 129 (105), Arizona 5, Utah 1, Nevada 1, California 11.

Mononucleosis: Connecticut 76, Ohio 2, Michigan 126, Minnesota 228, Maryland 31, South Carolina 5, Kentucky 12, Tennessee 29, Oklahoma 1, Montana 1, Idaho 24,

Oregon 7

Psittacosis: New York I (1), New Jersey I, Pennsylvania I, Michigan 4 (7), Alabama I, Washington I, California 12 (9).

Washington I, California 12 (1).

Purperal septicemia. New York 4 (2), Ohio 3, Florida 2 (2), Tennessee 3 (3), Mississippi 2 (8).

Lutkansas 5, Louisiana 2 (11), New Mexico 1 (4).

Rabies in animals: Maine 1, Rhode Island 2, New York 540 (646), Pennsylvania 75, Ohio 637 (761), Indiana 806 (723), Michigana 306 (313), Wisconsin 4, Minnesota 6 (1), Iowa 56, Missouri I, Kanasa 19 (62), Delaware I, Virginia 156, Week Minnesota 6 (1), Iowa 56, Missouri I, Kanasa 19 (62), Delaware I, Virginia 156, Week Alabama 370, (473), Arkansas 84 (63), Louisiana 90 (17), Oklahoma 131, Texas 1,271 (1971), Colorado 2 (13), New Mexico 2 (6), Arksona 47 (60), Utah I (7), California 280

Rat bite fever: Ohio 1.
Relation of the state of the scale of the scal

Yellow fever: Panama Canal Zone 5 fatal confirmed cases at a point outside the Canal Zone, about 22 miles from Panama City.

INCIDENCE OF DISEASE

No health department, State or local, can effectively prevent or control disease without knowledge of when, where, and under what conditions cases are occurring

UNITED STATES

REPORTS FROM STATES FOR WEEK ENDED JUNE 4, 1949

The incidence of poliomyelitis declined during the week for the country as a whole. A total of 139 cases was reported, as compared with 155 last week, 149 for the corresponding week last year, and a 5-year (1944-48) median of 71. The largest decrease was reported in Texas (72 last week to 37). Of 8 other States reporting currently more than 3 cases each, all except California (with a decline of 18 to 15) showed increases, the largest as follows (last week's figures in parentheses): Oklahoma 19 (7), South Dakota 7 (1), Tennessee 6 (0), Massachusetts 7 (2). A total of 835 cases has been reported since March 19 (average date of seasonal low incidence), as compared with 878 for the corresponding period last year and a 5-year median of 414.

The incidence of measles, 17,967 cases for the week, while lower than last week (19,834) and below the figures for the corresponding weeks of 1946 (26,347) and 1948 (23,883), is slightly above the 5-year median (16,130).

Of 19 cases of Rocky Mountain spotted fever reported currently (5-year median 23), 5 occurred in Virginia, 4 in Maryland, 2 each in North Carolina and Tennessee, 1 in Montana, and 1 each in 5 Middle Atlantic and East Central States. The total to date is 123. The largest number for a corresponding period of the past 5 years, 88, was reported in 1946 and in 1948.

Of 32 cases of tularemia (last week 20, 5-year median 20), the largest numbers occurred in Arkansas (11), Texas (8), and Georgia (3). The total to date is 530, 5-year median 392.

Of 52 cases of typhoid fever (5-year median 79), only Texas (with 9 cases) reported more than 3 cases.

During the week 1 case of psittacosis was reported in Michigan (Detroit).

Deaths recorded during the week in 94 large cities in the United States totaled 8,731, as compared with 9,008 last week, 8,606 and 9,192, respectively, in the corresponding weeks of 1948 and 1947, and a 3-year (1946-48) median of 9,192. The cumulative figure is 211,044, as compared with 215,021 for the same period last year. Infant deaths totaled 611, last week 661, 3-year median 661. The cumulative figure is 14,342, same period last year 15,098.

Telegraphic case reports from State health officers for week ended June 4, 1949

(Leaders indicate that no cases were reported)

(normale manage and comments and comments)	Enceph- alitis, enga meale menin- tious Measles gitis, monia myelitis spotted fever flows Measles goooccal flower flowe	33.5 10 1 16 1 16 1 16 1 1 16 1 1 16 1 1 16 1 1 1 16 1 1 1 1 10 1 1 1 1	5 (*) 2 1,483 7 161 5 1 490 2 1 490 2 1 183 2 1 1 83 3 1 1,698 3 59 1 1 140	3 2 2 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	2 3 3 1 6 2 4 4 1 1 6 1 7 7 3 3 7 7 7 3 3 7 7 7 3 3 7 7 7 3 3 7 7 7 3 3 7 7 7 3 3 7 7 7 3 3 7 7 7 3 3 7 7 7 3 3 7	2 214 006 2 74 (9) 2 8 8 14 006 2 1
	Diph- a t	(a)	40 M	D 03 03	- 8 -	8 - 49
	Division and State	Maine NEW ENGLAND New Hampshire Sermont Massechusetts Rhode Island Connecticut MINDER ATLANTIC	New York New Jorsey Pennsylvania Rast North Centrall Ohio	Michigan * Wisonain Wash North Chullal	Minnepota Joya Missouri North Dakota South Dakota Nebruska Kansa BOUTH ATLANTIC	Delaware Maryland Maryland District of Columbia, Viginia West Virginia Worth Oacolina South Carolina

• •	· - 5			5 0 5 0 0 0 0 0 1 1 0 0 0 0 0 0 0 0 0 0 0 0 0	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
8200	108	141 181	112	1,046 2,079	22, 476 (39tb) 0ct. 2 75, 130
9	-800	64		70	1, 180 (11th) Mar. 19 535 667
1 1 1 1	2018	1 1 1 1 1 ced 1	# 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	32 30	302
8 1 1 1 8 1 1 1 8 1 1 1 1 1 1 1		8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	6	4	342 218 (35th) Sept. 4 \$ 52 301
40 M	8100		800	1, 134 2, 458	53, 407 75, 274 (32nd) Aug. 14 76, 195 113, 845
-8			E 0 0 0 0 5 0	10	1123
@ m	225	0	15.13	130	11,757 (11th) Mar. 19 1834
7838	273		42	1, 234	46, 107
	8444		00	108	1, 782 3, 608 (37th) Sept. 18 2, 626 5, 112
8522	180 218 8218 833	881,360,93	386	17, 967	519, 301 443, 446 (35th) Sept. 4 571, 694 478, 392
119	1 400	- 7 8	8	923	71, 644 185, 879 (30th) July 31 107, 914 329, 915
		1	90	17 9	103
	8		es es es	117	3, 231 5, 567 (27th) July 10 8, 445 13, 123
Rantucky Temnessee Alabama Missisppi a. wrst bouth crytal	Arkansas Louisiana Oklahoma Texas Mountain	Montana Idaho Myoming Colorado New Mexico Articoa Utah -	PACIFIC Washington Oregon California	Total Median, 1944-48	Year to date, 22 weeks. Median, 1944-48. Beasonal low week ends. Since seasonal low week.

Period ended earlier than Saturday.
 Phe median of the 5 preceding corresponding periods; for pollomyelitis and typhoid fever the corresponding periods are 1944-45 to 1948-49, inclusive.
 New York City and Philadelphia only, respectively.
 Including cases reported as streptococcal infection and septic sore throat.
 Including partyphoid lever; reported septially as follows: New York I; Virginia I; Georgia I; Florida 2; Colorado I; Oregon I.
 Including partyphoid lever; reported septialed fever, Maryland, I case, April noise.
 Including partyphoid fever, Maryland, I case, April noise.
 Including party Poliomyelitis, South Carolina, week ended May 14, 1 case, Smallox-Diagnoses changed in certain States, weeks ended, as follows: Year 1949, South Dakota 1 case, Nay 1 1949.
 Patteorsi, Markingan I.
 Patteorsi, Markingan I.
 Ablates: Influenza 14; scalet lever I.
 Hawali Territory: Measles 89.

TERRITORIES AND POSSESSIONS

Puerto Rico

Notifiable diseases—4 weeks ended May 28, 1949.—During the 4 weeks ended May 28, 1949, cases of certain notifiable diseases were reported in Puerto Rico as follows:

Disease	Cases	Disease	Cases
Chickenpox Diphtheria Gonorrhea Influenza Malaria Measles	123 18 94 35 24 33	Syphilis. Tetanus. Tuberculosis (all forms) Typhold fever. Typhus fever (murine) Whooping cough	10 37 1

PLAGUE INFECTION IN SAN JUAN COUNTY, NEW MEXICO, AND SEVIER COUNTY, UTAH

Under date of June 1, plague infection was reported proved in a pool of 24 fleas from 20 grasshopper mice, Onychomys leucogaster, trapped, on May 17, 20 to 23 miles southeast of Bloomfield, San Juan County, N. Mex., along U. S. Highway 44, and in a pool of 298 fleas from 78 ground squirrels, Citellus armatus, shot May 12 in Fish Lake National Forest 33 miles southeast of Richfield on Fish Lake Road, Sevier County, Utah.

DEATHS DURING WEEK ENDED MAY 28, 1949

[From the Weekly Mortality Index, issued by the National Office of Vital Statistics]

	Week ended May 28, 1949	
Data for 94 large cities of the United States:		
Total deaths	9, 008 8, 324	9, 013
Median for 3 prior years	202, 313	206, 415
Deaths under 1 year of age	661	675
Median for 3 prior years	675	
Deaths under 1 year of age, first 21 weeks of year	13, 731	14, 437
Policies in force	70, 393, 900	71, 072, 486
Number of death claims	13, 287	12, 697
Death claims per 1,000 policies in force, annual rate	9.8	9. 3
Death claims per 1,000 policies, first 21 weeks of year, annual rate	9.7	10. 2

FOREIGN REPORTS

CANADA

Provinces-Notifiable diseases-Week ended May 14, 1949.—During the week ended May 14, 1949, cases of certain notifiable diseases were reported by the Dominion Bureau of Statistics of Canada as follows:

Disease	Prince Edward Island	Nova Scotia	New Bruns- wick	Que- bec	On- tario	Mani- toba	Sas- katch- ewan	Alber-	British Colum- bia	Total
Chickenpox		27	3	239	546	31	40	46	115	1,050
Dysentery, bacillary	*******						1	1	******	2
Encephalitis, infectious German measles Influenza	*******	6 17		317	64	25	122	40	7	581
Measles	*********	50	4	268	278	193	207	349	351	1, 700
cal		17		50	323	16	*3	8	48	46
Poliomyelitis		1 3							******	157
Scarlet fever		14	8	62	63 28	29	8	15 13	11 62	225
phoid feverUndulant fever			3	4	1	******		1	*******	1
Venereal diseases:						-		-		001
Gonorrhea	1 2	7 6	12	124 77	61	27	13	28	65	338
Whooping cough	2	1	3	63	22		1			87

WORLD DISTRIBUTION OF CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER

From consular reports, international health organizations, medical officers of the Public Health Service, and other sources. The reports contained in the following tables must not be considered as complete or final as regards either the list of countries included or the figures for the particular countries for which reports are given.

CHOLERA

(Cases)

Note.—Since many of the figures in the following tables are from weekly reports, the accumulated total⁸ are for approximate dates.

	January-	April	Mı	ay 1949—	week end	led—
Place	March 1949	April 1949	7	14	21	28
ASIA	18					
Burma Bassein Moulmein	20 4	6	7 7	1	56 56	1
India Allahabad Bombay	27, 968 1	10, 889	1 1, 523	1 871	184	40
Calcutta	3 1, 791 11	3 1, 453 15	3 238 4	3 212 5	3 173 6	4
Lucknow	9 10	9 15 6	5 9 1	10	8	18
Raj Samand Tuticorin India (French):	14				1	9
Fondicherry	100					

See footnotes at end of table.

CHOLERA-Continued

No.	January- March	January- April		May 1949—week ended—				
Place	1949	April 1949	7	14	21	28		
ASIA—continued Indochina (French): Annam Cambodia Cochinchina Pakistan Chittagong Dacca Lahore Sian Bangkok	50 11 1 10,640 42	3 21 2 2,470 13 45	24 3 91	2 11 5 6 6 6	1			

³ Includes imported cases. ¹ Preliminary figures. ² Imported.

PLAGUE

(Cases)

	1		1	1	1	1
AFRICA						
Basutoland	12		1		1.	
	4	2				
Belgian Congo	4	2	*****			
	- 1	- 2	******			
British East Africa:				11500		1
Kenya	1					
Tanganyika	15	********				
Madagascar	48	12		12		
Tananarivo	2					
Rhodesia, Northern	2					
Union of South Africa	26					0.000000
Union of podem zenions account to the contract	-					
ASIA		-				
Burma	1 329	16			5	
Mandalay	1					
Moulmein	232 1 50	4			110314	
Rangoon	19	19			31	
Chi-					- 4	******
China:						
Chekiang Province	5	2	******			*******
Wenchow	. 5	2				
Fukien Province	20	*********				*******
Klangsi Province	9					
India	19, 202	3, 992	4 23	4 20	4 37	100000000000000000000000000000000000000
Indochina (French)	30	6 31	1	81		
Annam	00	7	i	81		
Cambodia	17	3			*******	********
Carlondia		6 21	*******	******		******
Cochinchina	7	* 21	******	******		
Laos	3	*********			*******	
Java	2	3	******		******	
Siam	136	12				
						-
EUROPE						
Portugal: Azores	3					
SOUTH AMERICA		-				
Peru:		100				
Lambayeque Department	7	********				
Piura Department	3					
Venezuela:						
Aragua State	Constitution of	1	134. E			
· versibility or and a second						
OCEANIA		-			15.75	
OUBANIA					1 2	
						DV Day
Hawaii Territory: Plague infected rats						

¹ May 1-10, 1949.
2 Includes imported cases.
3 Imported.
4 Reports from Calcutta and Cawnpore only.
5 Pneumonic plague.
6 Includes cases of pneumonic plague.
7 Plague infection has been reported in Hawaii Territory as follows: On March 12, 1949, in a mass inoculation of 2 pools of tissue from 10 rats (8 and 2), taken on Maui Island; on March 16, 1949, in mass inoculation of 3 pools of 29 fleas (7, 12, and 10), from rats trapped on the Island of Hawaii.

SMALLPOX

(Cases)

(P=present)

Place	January- March 1949	April 1949	May 1949—week ended—			
			7	14	21	28
AYRICA						
Iperia	87	11		1.5	******	
Ingola	118	1 199		******	******	*****
Belgian Congo	2 543	s 188		*******		
British East Africa:	9	- 4				
Kenya Nyasaland	542	146	11	30	******	
Tanganyika	122		******	******	*******	*****
TanganyikaUganda	30	2		18	******	*****
ameroon (French)	13	6 34	*******	1 10	86	
)ahomev	152	2	*******	- 10		*****
Sypt. Sthiopis French Equatorial Africa	1					
Propen Equatorial Africa	4	2		18	*****	
French Guinea	1			******	******	*****
French Guinea French West Africa: Haute Volta	43	14		12	******	*****
Pambia	5	35	******	11	******	
Vory Coast	86	83	******	ii	*******	*****
dorocco	53	1 3	******		*******	
Mozambique	1.149	435				*****
Nigeria Niger Territory	1, 149 119	133		1 49		
Portuguese Guines	1				******	
Portuguese GuineaRhodesfa:	0 9					
Northern	4		******	******	******	
Southern	135	**********	******	******	******	*****
Serra Leone	14 76	1 4	*******		******	*****
Slerra Leono	28	i	8	14		
nden (French)	28 82	54				
Pogo (French)	46	11		15	81	
udan (Anglo-Egyptian) udan (French) Togo (French) Union of South Africa	124	P	Im			
ASIA	5 15	13.				
Afghanistan	15					
Arabia	30	14	******	******		*****
Bahrein Islands Burma	760	8 55	8 12	8 15	8 12	
China	4 651	4 96	4.11	4 17		
India	22, 083	13, 607	1, 233	724		
ndia (French)	1					
India (French) India (Portuguese) Indochina (French)	124	23		25	3	*****
Indochina (French)	1, 950	166	17		1000	LI BU
ran	156 245	18	1 7	1 5	48	
IraqIsrael	2	10				
Japan	16	23	8	******		
ava	1, 993	1, 174	6 150	6 246	4 297	8
Korea	544	********				****
T - borners	112	9	1	5		*****
Malay States (Federated)	42	302		******		
Malay States (Federated) Pakistan Philippine Islands: Mindero Island Tablas Island Portuguese Timer	1, 880	002	******	*******	*******	
Mindore Island	2					
Tables Jeland		2			******	
Portuguese Timor	4			******		
	37					
Straits Settlements: Singapore	12				7	*****
Sumatra	4 36	13	*1	23	19	1000
Syria Fransjordan	221	41 51	8	7	4	1 39
Purkey. (See Turkey in Europe.)			1	120	1000	100
EUROPE		1	1	1	1	1
Great Britain:		7 11	1	2	2016	1
England and Wales	9	. 11				
ItalyPortugal: Lisbon	2 3	********				
Spain.		1				
Turkey	83	4	******	1	******	
	1 5 7 1					
NORTH AMERICA	2			-		
Cuba: Habana Guatemala	1 7					
Mexico.		7	1	-		1

See footnotes at end of table.

SM ALLPOX-Continued

Place	January- March 1949	April 1949	May 1949—week ended—				
			7	14	21	28	
SOUTH AMERICA							
rgentina		3 8 54	******	******	91		
Prazil	² 55 556	26			.1		
olombia	000						
cuador	2 348	2 59					
araguay	*1						
eru	387	298					
Venezuela.	2 522	19		9 28			

TYPHUS FEVER*

(Cases)

(P=present)

Algeria	1.	- prosens/				460	
Algeria 27 12 13	APPROA						
Basutoland		97	12		13		
Belgian Congo.					-0	*******	******
Egypt			2 10				
Eritrea							
Sthiopia	Pritron			******			
Abya			-				
Madagasoar: Tananarive			33	9			
Morocco. S 32 32 72 72 73 74 74 74 74 74 74 74			1 00	-			
Cunisia			1				
Indicate							
ASIA Asia				p		*******	
Afghanistan 1, 191 1 1 1 1 1 1 1 1 1	mon of South Africa	140	1	-	******	********	
A						1	1
Ceylon: Colombo							
China			1				
India (Portuguese)							
Addition Addition	hina		2				
1				41			******
Page			2				
13 8 3 1 1 1 1 1 1 1 1 1	ndochina (French)				2		
Section Sect	ran	62	48				
Span	agp	13	8	3	1	1	
125	apan	# 60	15				
Sakistan		125					
Palestine 100 1 1 1 1 1 1 1 1	ebanon	11					
alestine 100	akistan	320	180	9	11	19	
hilippine Islands: Manila		100					
traits Settlements: Singapore	hilippine Islands: Manila	1					
yria. 2 2 6 5 ransjordan 11 31 6 1 1 urkey. (See Turkey in Europe.) 1 1 31 6 1 1 elgium. 1 71 94 31 31 31 31 31 31 31 32 3	traits Settlements: Singapore	*1				1	
Transjordan			2	2	6	5	
EUROPE elgium algaria. zechoslovakia rance reat Britain: Island of Malta 2 tungary aly Sicily oland 1 1 2 2 1 1 2 2 1 1 2 1 1	ransjordan	11	31		i	1	
elgium	urkey. (See Turkey in Europe.)	10.7			1		
Salgaria	EUROPE		100	1990		T. I	
Zechoslovakia							
zechoslovakia 2 6 1 rance 2 1 2 reat Britain: Island of Malta 1 2 1 reece 221 21 2 2 ungary 12 3 1 1 aly 27 3 1 1 Sicily 13 5 29 ortugal: Lisbon 2 2 2 umanis 297 297	ulgaria	71	94	31			
rance. reat Britain: Island of Malta 3 reece. 21 21 21 22 22 23 21 21 21 21 21 22 27 27 28 28 29 29 20 20 20 20 20 20 20 20 20 20 20 20 20		2	6	1			******
reece		2					
reece	reat Britain: Island of Malta 3	1	2				
Ungary		3 21		12	12		
aly 27			3		1	1	
Sicily							
oland 150 29							
ortugal: Lisbon 2 2 umania. 297	oland		20				
umania 297							
pain.		1		1			*****
urkey	nrkov		91	1	1	9	
Ugoslavia. 82 19 1 10	nonelavia						1

See footnotes at end of table.

¹ May 1-10, 1949. ² Includes alastrim. ³ May 11-20, 1949. ⁴ Includes imported cases. ⁵ In Moulmein and Rangoon only. ⁶ In Batavia only. ⁷ Imported. ⁸ January 1-April 30, 1949. ⁹ Alastrim.

TYPHUS FEVER-Continued


Place	January- March 1949	April 1949	May 1049—week ended—			
			7	14	21	28
Costa Rica 3 Cuba 3 Guatemala Jamaica 3 Mexico 3 Puerto Rico 3	8 1 7 5 46 5	5 1 1 11 2	6		1	
Bolivia. SOUTH AMERICA Brazil Chile Colombia Curacao ² Ecuador ³ Peru Venezuela ³	8 2 43 574 2 84 5 8	31 34 1 23 2 7	9	2	1	
Australia ³	33	17	1	******		

YELLOW FEVER

(C—cases; D—deaths)				
Belgian Congo: Stanleyville Province	5			
Nigeria: C	32			
Panama: NORTH AMERICA Pacora	48			
Brazil: Amazonas State D Para State D	1			

¹ May 1-10, 1949. ² Murine type. ³ Includes murine type. ⁴ Imported. ⁶ Corrected figure. ⁶ Type unspecified.

Between Sekondi and Cape Coast.
 Reported May 2, 1949.
 Cases admitted to Lagos Hospital from ship that arrived from two other ports in Nigeria—Warri and Burutu.
 Reported January 15, 1949.
 Date of occurrence November 11-December 30, 1948.
 Five cases, all fatal, confirmed; 3 suspected cases.
 Reported April 6, 1949, at Acara.



+++

The Public Health Reports is printed with the approval of the Bureau of the Budget as required by Rule 42 of the Joint Committee on Printing.

The Public Health Reports, first published in 1878 under authority of an act of Congress of April 29 of that year, is issued weekly by the Public Health Service through the Division of Public Health Methods, pursuant to the following authority of law: United States Code, title 42, sections 241, 245, 247; title 44, section 220.

It contains (1) current information regarding the incidence and geographic distribution of communicable diseases in the United States, insofar as data are obtainable, and of cholera, plague, smallpox, typhus fever, yellow fever, and other important communicable diseases throughout the world; (2) articles relating to the cause, prevention, and control of disease; (3) other pertinent information regarding sanitation and the conservation of the public health.

The Public Health Reports is published primarily for distribution, in accordance with the law, to health officers, members of boards or departments of health, and other persons directly or indirectly engaged in public health work. Articles of special interest are issued as reprints or as supplements, in which forms they are made available for more economical and general distribution.


Requests for and communications regarding the Public Health Reports, reprints, or supplements should be addressed to the Surgeon General, Public Health Service, Washington 25, D. C. Subscribers should remit direct to the Superintendent of Documents, Washington 25, D. C.

Librarians and others should preserve their copies for binding, as the Public Health Service is unable to supply the general demand for bound copies. Indexes will be supplied upon request.


+++


UNITED STATES GOVERNMENT PRINTING OFFICE, WASHINGTON, D. C. 1 1949

For sale by the Superintendent of Documents, U. S. Government Printing Office, Washington 25, D. C. Price 10 cents, Subscription price \$4.00 a year.



PUBLIC LIBRARY
AUG 201949
DETROIT





Index, v. 64, part 1

Public Health Reports

Issued Weekly by the
PUBLIC HEALTH SERVICE

Index
Volume 64—Part I
Nos. 1-25
January-June 1949



FEDERAL SECURITY AGENCY

PUBLIC HEALTH SERVICE

FEDERAL SECURITY AGENCY

Oscar R. Ewing, Administrator

PUBLIC HEALTH SERVICE

Leonard A. Scheele, Surgeon General

Division of Public Health Methods G. St. J. Perrott, Chief of Division

The printing of this publication has been approved by the Director of the Bureau of the Budget (August 10, 1949).

The Public Health Reports, first published in 1878 under authority of an act of Congress of April 29 of that year, is issued weekly by the Public Health Service through the Division of Public Health Methods, pursuant to the following authority of law: United States Code, title 42, sections 241, 245, 247; title 44, section 220.

It contains (1) current information regarding the incidence and geographic distribution of communicable disease in the United States, insofar as data are obtainable, and of cholera, plague, smallpox, typhus fever, yellow fever, and other important communicable diseases throughout the world; (2) articles relating to the cause, prevention, and control of disease; (3) other pertinent information regarding sanitation and the conservation of the public health.

The Public Health Reports is published primarily for distribution, in accordance with the law, to health officers, members of boards or departments of health, and other persons directly or indirectly engaged in public health work. Articles of special interest are issued as reprints or as supplements in which forms they are made available for more economical and general distribution.

Requests for and communications regarding the Public Health Reports, reprints, or supplements should be addressed to the Surgeon General, Public Health Service, Washington 25, D. C. Subscribers should remit direct to the Superintendent of Documents, Washington 25, D. C.

Librarians and others should preserve their copies for binding, as the Public Health Service is unable to supply the general demand for bound copies. Indexes will be supplied upon request.

UNITED STATES GOVERNMENT PRINTING OFFICE, WASHINGTON: 1950

For sale by the Superintendent of Documents, U. S. Government Printing Office Washington 25, D. C. Index, 10 cents. Regular issues, 10 cents. Subscription price \$4.00 a year.

Public Health Reports Subject Index

Key to Dates and Pages

No.	Date of issue	Pages	No.	Date of issue	Pages
1	Jan. 7	1-40	14	Apr. 8	439-469
2	Jan. 14	41-68	15	Apr. 15	471-498
3	Jan. 21	69-92	16	Apr. 22	499-522
4	Jan. 28	93-120	17	Apr. 29	523-550
5	Feb. 4	121-160	18	May 6	551-588
6	Feb. 11	161-200	19	May 13	589-620
7	Feb. 18	201-228	20	May 20	621-656
8	Feb. 25	229-258	21	May 27	657-684
9	Mar. 4	259-302	22	June 3	685-710
10	Mar. 11	303-329	23	June 10	717-748
11	Mar. 18	331-354	24	June 17	749-784
12	Mar. 25	355-401	25	June 24	785-81
13	Apr. 1	403-438			

Page Accident facts from accident fatalities [Halpin]388 Accident prevention: Introduction [Armstrong]_____ Research [King]____ A State health department's responsibility [Beelman] _____363 Accident-prone individual [Alexander]___ Accidents, the home environment and [Board]_____383 Actinomycosis cases: Annual report for 1948 by State......806 Quarterly reports by State_____ Alameda County, California, nursing service..... Alaska: Communicable diseases _____35, 65, 113, 155, 196, 219, 247, 297, 326, 350, 393, 435, 463, 495, 541, 585, 617, 645, 677, 711, 743, 809

American Board of Preventive Medicine and Public Health, Inc., formation_512

Animal colony, experimental, in tropical West Africa [Poindexter]_____57

Influenza_

Anthrax cases:	Page
Annual report for 1948 by State	_802
Quarterly reports by State224	, 652
Weekly State reports65, 89, 155, 219, 247, 297, 350, 495, 585, 617, 677, 743	519,
Antibiotic-producing coliform organisms, prevalence of [Halbert and Gravatt	
Aureomycin: See Pertussis and	
Australia: Measles	940
Poliomyelitis	-114
В	
Blastomycin: See Histoplasmin and	
Board of Preventive Medicine and Public Health, American	_512
Botulism cases:	
Annual report for 1948 by State	
Quarterly reports by State228	, 656
British National Health Service [Davis]	_161
Brucella abortus, isolation from hogs [McCullough, Eisele and Pavelchek]	537
C	
Canada:	
Communicable diseases	36,
66, 90, 114, 156, 198, 220, 249, 299, 327, 352, 394, 436, 497, 542, 586, 618, 648, 680, 714, 745, 782,	520, 811
Diphtheria	220
Cancer:	
Cases:	
Annual report for 1948 by State	806
Quarterly reports by State228,	
National Cancer Institute program of postgraduate training for physi-	
cians [Spencer]	750
Professional education for, control [Deibert]	749
Teaching in dental schools [Kaiser]	764
Teaching in medical schools [Kaiser]	757
Tools for professional cancer education, new	769
Ceylon: Poliomyelitis 91,	352
Chickenpox cases:	
Annual report for 1948 by State	802
Quarterly reports by State 224,	652
Cholera: Foreign reports	91,
115, 199, 222, 252, 301, 328, 353, 395, 521, 544, 587, 619, 649, 681, 715, 746,	467, 811
Choriomeningitis, lymphocytic, cases:	
Annual report for 1948 by State	806
Quarterly reports by State 228,	656
Cobalt and the dust environment of the cemented tungsten carbide industry	
[Fairhall, Keenan and Brinton]	485
Considinidom vensis esses	
	806
Quarterly reports by State228,	656
	919

Collodion agglutination:	· P	agi
Technique		290
Test for histoplasmosis [Saslaw and Campbell]	4	124
Colombia: Typhoid fever	4	136
Colorado tick fever cases:		
Annual report for 1948 by State	8	806
Quarterly reports by State	2	228
Commissioned Corps, Regular, Public Health Service examinatio engineer officers		159
Communicable disease charts:		
Diphtheria	11, 5	17
Measles 111, 245, 391,	517, 646, 7	80
Meningitis, meningococcal	111, 3	91
Poliomyelitis		
Scarlet fever		
Typhoid and paratyphoid fever		
Whooping cough		
Communicable diseases—Current incidence by State	1	33.
63, 87, 110, 153, 194, 217, 244, 295, 324, 348, 390, 4		
516, 539, 583, 615, 643, 675, 709,		
Cunjunctivitis cases:	141, 111, 0	,,,,
Annual report for 1948 by State	9	302
Quarterly reports by State		
Cuba		
Habana—Communicable diseases91, 221, 436,	485 748 7	122
Provinces—Notifiable diseases 91, 221, 437,		
Trovinces—Avoinable diseases	100, 110, 1	00
D		
DDT lastice offerts or demotis note under colons and field and	41	
DDT dusting, effects on domestic rats under colony and field condi		00
[Dent, Morlan and Hill]	0	66
		20
Large cities—Weekly reports (Weekly mortality index)		38,
68, 92, 120, 158, 197, 223, 258, 298, 329, 354, 401, 4		
515, 550, 588, 620, 651, 674, 713, 7		
Tuberculosis—Map		08
Dengue cases:	0	00
Annual report for 1948 by State		06
Quarterly reports by State	228, 6	50
Dermatitis cases:		
Annual report for 1948 by State		06
Quarterly reports by State	228, 6	56
Dextrose serum tellurite plating medium. Its use in diphtheria diag		
[Whitley and Damon]	20	01
Diarrhea cases:		
Annual report for 1948 by State	80	06
Quarterly reports by State	228, 68	56
Diarrheal disease control studies. II. Conical net for collecting	flies	
[Major and How]	6.6	1.4

Diphtheria:	Page
Canada	220
Cases:	
Annual report for 1948 by State	802
Quarterly reports by State224	, 652
Weekly State reports	34,
64, 88, 112, 154, 195, 218, 246, 296, 325, 349, 392, 434, 462,	494,
518, 540, 584, 616, 644, 676, 710, 742, 778,	, 808
Diagnosis	457
Dog bite cases:	
Annual report for 1948 by State	806
Quarterly reports by State228	, 656
Drugs, antimalarial, summary [Cooper]	717
Dust environment	485
Dysentery cases:	
Annual report for 1948 by State	802
Quarterly reports by State224	652
4	
- E	
Education, cancer 749, 750, 757, 764.	760
Encephalitis cases: Infectious:	709
Annual report for 1948 by State	802
Quarterly reports by State 224,	
Weekly State reports	34.
64, 88, 112, 154, 195, 218, 246, 296, 325, 349, 392, 434, 462,	
518, 540, 584, 616, 644, 676, 710, 742, 778, Other forms:	
Annual report for 1948 by State	806
Quarterly reports by State 228.	
Engineer officers, Regular Corps, Public Health Service examinations	159
Epidemiological terms, suggested. "Infection unit" and "index of aggre-	
gation" [Forsbeek]	343
Erysipelas cases: Annual report for 1948 by State	006
	806
Quarterly reports by State	
engineer officers	159
· · ·	
and the second s	
Favus cases: Annual report for 1948 by State	806
Quarterly reports by State	228
Filariasis cases: Annual report for 1948 by StateFinland;	806
Influenza	251
Notifiable diseases249, 327, 394, 649,	714
Flocculation test as a possible method for differentiating immunologic types of the poliomyelitis virus [Roberts]	212
Fluoroscopic screens, persistence of [Van Allen]	560

	-
Food poisoning cases:	Page
Annual report for 1948 by State	806
Quarterly reports by State228	
Foreign reports—Quarantinable diseases	36,
66, 90, 114, 156, 198, 220, 249, 299, 327, 352, 394, 436, 465, 520, 542, 586, 618, 648, 680, 714, 745, 782	
G	
German measles cases:	
Annual report for 1948 by State	802
Quarterly reports by State224	
Gold Coast: Meningitis, cerebrospinal 437, 465, 542	
Gonorrheal specimens, simple and efficient transport method for [Peizer,	, 010
Steffen and Klein]	599
Grants-in-aid for public health, ten years of—Review of Bulletin 300	000
[Mountin, Hankla and Druzina]	490
Granuloma inguinale cases:	200
Annual report for 1948 by State	806
Quarterly reports by State228	
quarterly reports by booteness	, 000
. Н	
Hawaii:	
Communicable diseases	35,
65, 89, 113, 155, 196, 219, 247, 297, 326, 350, 393, 435, 463,	495,
519, 541, 585, 617, 645, 677, 711, 743, 779,	809
Plague (in rodents and fleas)	464
Heart disease, statistical studies of:	
III. Heart disease associated with other major causes of death as pri-	
mary or contributory cause [Gover]	104
IV. Mortality from heart disease (all forms) related to geographic	
section and size of city [Gover]	439
Hepatitis: Japan	250
Histoplasmin and blastomycin, a comparison between, by the collodion	
agglutination technique. [Saslaw and Campbell]	290
Histoplasmosis:	
Collodion agglutination test for [Saslaw and Campbell]	424
Use of yeast-phase antigens in a complement fixation test. III.	
Preliminary results with human sera. [Campbell and Saslaw]	551
Hookworm disease cases:	
Annual report for 1948 by State	802
Quarterly reports by State224,	652
Hospital maintenance and operation, State legislation for minimum stand-	
ards of [Hoge and Steinle]	69
T	
Impetigo contagiosa cases: Annual report for 1948 by State	000
	806
Quarterly reports by State228, Industrial hygiene agencies, work of State and local [Trasko]	
	471
Infant mortality and socioeconomic factors, relationship between, in urban	991
areas [Altenderfer and Crowther]	331
Index of aggregation, epidemiological term suggested	343
Infection unit, epidemiological term suggested	343

Influenza:	Page
Alaska	192
Cases:	
Annual report for 1948 by State	802
Quarterly reports by State 224	, 652
Weekly State reports 34, 64, 88, 112, 154, 195, 218, 246, 296, 325,	349,
392, 434, 462, 494, 518, 540, 584, 616, 644, 676, 710, 742, 778	
Europe 66, 157, 193	
Austrian Tyrol	193
Belgium	251
England	157
Finland	251
France 157, 193,	-
	251
Germany	
Holland 157,	
Italy66, 157,	
Switzerland 157,	
Turkey	157
New Zealand	
Nicobar Islands (India)	543
Puerto Rico	193
Report on recent outbreaks 157, 192,	251
United States	192
Venezuela	193
WHO report	251
Interrelation of social problems (tuberculous patients)—editorial [Anderson]_	685
J	
Jamaica: Notifiable diseases 36, 198, 300, 465, 586,	714
Japan:	
Hepatitis	250
Notifiable diseases 90, 221, 352, 520, 648,	
Jaundice cases:	104
Annual report for 1948 by State	806
Quarterly reports by State228,	
Quarterly reports by State	000
\mathbf{v}	
K	
Kala-azar: Annual report for 1948 by State	000
Kala-azar: Annual report for 1948 by State	806
L	
	228
Leprosy cases:	
	806
Quarterly reports by State228,	
	201
Loeffler's medium, replacement for	457
Louisville, Ky., murine typhus fever in [Good and Kotcher]	229
Lymphogranuloma venereum cases:	
Annual report for 1948 by State	806
Quarterly reports by State 228	656

M	Page
Madagascar: Notifiable diseases 37, 156, 250, 467, 5 Malaria:	43, 715
Cases:	
Annual report for 1948 by State	802
Quarterly reports by State2	
Drugs	717
Map—White and nonwhite tuberculosis deaths as a percent of all tuber- culosis deaths for each State and for the United States, 1947	
Mauritius: Poliomyelitis66, 250, 3	52, 497
Measles:	
Australia	294
Cases:	
Annual report for 1948 by State	
Quarterly reports by State2	24, 652
Weekly State reports	34,
64, 88, 112, 154, 195, 218, 246, 296, 325, 349, 392, 434, 46 518, 540, 584, 616, 644, 676, 710, 742, 7	
Meningitis:	
Cerebrospinal:	day.
Gold Coast 437, 465, 5	
Nigeria	618
Meningococcal:	
Annual report for 1948 by State	802
Quarterly reports by State2	
Weekly State reports	34,
64, 88, 112, 154, 195, 218, 246, 296, 325, 349, 392, 434, 46	
518, 540, 584, 616, 644, 676, 710, 742, 7	18, 808
	303
Act, national Clinic: Community [Davens and Lemkau]	657
First annual report of Prince Georges County clinic [Ross]	797
Metopon hydrochloride. An experiment in clinical evaluation [Eddy]	93
Milk sanitation ratings of 90 percent or more during 1947 and 1948	241
Mononucleosis cases:	211
Annual report for 1948 by State.	806
Quarterly reports by State2	1000
M1:324	20, 000
Annual report for 1948 by State	802
Quarterly reports by State	
Weekly State reports	33,
63, 87, 110, 153, 194, 217, 244, 295, 324, 348, 390, 433, 46: 516, 539, 583, 615, 643, 675, 709, 741, 77	1, 493,
Morphine derivative, new: Metopon hydrochloride	93
Mortality:	
Heart disease 10	
Infant, and socioeconomic factors 403, 40	331
Tuberculosis 403, 40 Mumps cases:	0, 087
Annual report for 1948 by State	802
Quarterly reports by State	
	0, 200
800504-50-2	

N

	Page
National Health Service, British [Davis]	161
National Mental Health Act, how it works [Lowry]	303
Netherlands: Psittacosis 157	, 198
New Zealand:	
Influenza	, 783
Notifiable diseases 251, 300, 466, 542	
Poliomyelitis	, 327
Nicobar Islands (India): Influenza	543
Nigeria: Meningitis, cerebrospinal	618
Norway: Notifiable diseases	, 680
Notifiable diseases:	
Fourth quarter, 1948	652
Third quarter, 1948	224
Year 1948	802
Nursing service: Alameda County (Calif.)	523
0	
Ophthalmia cases:	
Annual report for 1948 by State	802
Quarterly reports by State224	, 652
P	
Panama Canal Zone: Notifiable diseases 248, 351, 647, 679	
Panama carrier (typhoid fever)	671
Paratyphoid fever cases:	
Annual report for 1948 by State	804
Quarterly reports by State	654
Pasteurella pestis culture without transfer, twenty-five year survival of	
[Francis]	238
Pellagra cases:	
Annual report for 1948 by State	802
Quarterly reports by State224	
Pertussis and aureomycin [Bell, Pittman and Olson]	589
Philippines:	
Annual report of the tuberculosis control division [Young]	123
Public health rehabilitation program [Young]	123
Plague:	
Culture survival without transfer [Francis]	238
Diagnosis procedures, effect of sodium fluoroacetate (1080) in poisoned	
rats on. Preliminary report [Gratch, Purlia, and Martin]	339
Foreign reports	38,
67, 92, 116, 157, 199, 222, 253, 301, 328, 353, 396, 437, 467,	
521, 545, 587, 619, 649, 681, 715, 746,	812
Infection:	-
Arizona	
Colorado	781
New Mexico	
Texas	
Utah	
Washington 36 208 464 406 670	712

Pneumonectomy followed by immediate thoracoplasty [Iverson and Skinner]	Page
Pneumonia (all forms) cases:	
Annual report for 1948 by State	802
Quarterly reports by State224,	652
Weekly State reports	34,
64, 88, 112, 154, 195, 218, 246, 296, 325, 349, 392, 434, 462,	494,
518, 540, 584, 616, 644, 676, 710, 742, 778,	
Poliomyelitis:	
Australia	114
Cases:	
Annual report for 1948 by State	804
Quarterly reports by State226,	654
Weekly State reports	34,
64, 88, 112, 154, 195, 218, 246, 296, 325, 349, 392, 434, 462,	494,
518, 540, 584, 616, 644, 676, 710, 742, 778,	
Ceylon	91
Mauritius	497
New Zealand	114
Prevalence of, in 1948 [Dauer]	733
Reunion Island	544
Virus, a flocculation test as a possible method for differentiating im-	
munologic types of [Roberts]	212
Prince Georges County, Maryland, pilot mental health clinic	797
Psittacosis:	
Cases:	
	806
Quarterly reports by State228,	
Weekly State reports	
Netherlands 157,	
Public health nursing service, time analysis of. Alameda County Health	100
Department, 1946-47 [Malcolm, Moreland, Hopkins and Snyder]	523
Public Health Service:	
	159 608
Puerperal septicemia cases:	
Annual report for 1948 by State	806
Quarterly reports by State	656
Puerto Rico:	
Influenza	193
Notifiable diseases 197, 248, 351, 744,	810
Q	
Q fever:	
Cases:	
	806
	228
	541
Studies in southern California:	
II. An epidemiological study of 300 cases [Beck, Bell, Shaw and	
Huebner]	41
III. Effects of pasteurization on survival of C. burneti in naturally	
infected milk [Huebner, Jellison, Beck and Wilcox]4	199

R	Page
Rabies cases:	Page
In animals:	
Annual report for 1948 by State	806
Quarterly reports by State 228	. 256
Weekly State reports	
88, 112, 154, 195, 218, 246, 296, 325, 349, 392, 434, 462, 494,	
540, 584, 616, 644, 676, 710, 742, 778,	
In man:	
Annual report for 1948 by State	804
Quarterly reports by State 226	654
Weekly State reports	113
Rabies problems and control. A nation-wide program [Steele and Tierkel]	785
Raffinose serum tellurite agar slants as a replacement for Loeffler's medium	
in diphtheria diagnosis [Whitley and Damon]	457
Rat bite fever cases:	
Annual report for 1948 by State	806
Quarterly reports by State	656
Relapsing fever cases:	-
Annual report for 1948 by State	806
Quarterly reports by State 228,	
Reunion Island: Poliomyelitis	544
Rheumatic fever cases:	
Annual report for 1948 by State	804
Quarterly reports by State226,	
Rickettsialpox cases:	
Annual report for 1948 by State	806
Quarterly reports by State	
Ringworm cases:	
Annual report for 1948 by State	806
Quarterly reports by State228,	
Rocky Mountain spotted fever cases:	-
Annual report for 1948 by State	804
Quarterly reports by State226,	654
Weekly State reports34	
88, 112, 154, 195, 218, 246, 296, 325, 349, 392, 434, 462, 494,	
540, 584, 616, 644, 676, 710, 742, 778,	
8	
Salmonella typhosa 58, avirulent isolate of (Panama carrier) [Batson,	
Landy and Abrams]	671
Scabies cases:	
Annual report for 1948 by State	806
Quarterly reports by State228,	656
Scarlet fever cases:	
Annual report for 1948 by State	804
Quarterly reports by State226,	654
Weekly State reports 34,	64,
88, 112, 154, 195, 218, 246, 296, 325, 349, 392, 434, 462, 494,	518,
540, 584, 616, 644, 676, 710, 742, 778,	

The state of the s	
Schistosomiasis cases:	Page
Annual report for 1948 by State	806
Quarterly reports by State22	3, 656
Screens, fluoroscopic	560
Septic sore throat cases:	
Annual report for 1948 by State	804
Quarterly reports by State 220	3, 654
Sickness absenteeism among industrial workers. Third and fourth quarters of 1948 [Gafafer]	641
Silicosis cases:	-
Annual report for 1948 by State	806
Quarterly reports by State 228	
Smallpox:	
Cases:	
Annual report for 1948 by State	804
Quarterly reports by State226	6, 654
Weekly State reports 3	
88, 112, 154, 195, 218, 246, 296, 325, 349, 392, 434, 462, 494, 540, 584, 616, 644, 676, 710, 742, 778	518,
Foreign reports 3	
92, 117, 157, 199, 222, 254, 301, 328, 353, 397, 438, 468, 498	
546, 587, 619, 649, 682, 716, 747, 784	
Vaccination, effect of, on the outcome of pregnancy [Bellows, Hyman	, 010
and Merritt]	319
Social problems of tuberculous patients.	685
Socioeconomic factors and infant mortality, relationship between, in urban	000
areas [Altenderfer and Crowther]	331
Sodium fluoroacetate (1080) in poisoned rats, effect of, on plague diagnosis	001
procedures [Gratch, Purlia and Martin]	339
Sodium hydroxide concentration techniques in tuberculosis diagnosis	574
Specimens, gonorrheal, transporting of	599
State health departments, current organizational pattern of statistical	
activities in [Swinney]Studies of patients discharged from tuberculosis sanatoria. II. Mortality	621
rates associated with selected characteristics of the patient population	007
[Brewster and Fletcher]Switzerland:	687
Influenza157	102
	466
Notifiable diseases	400
T	
Tetanus cases:	
Annual report for 1948 by State	804
Quarterly reports by State 226,	654
Thoracoplasty following pneumonectomy [Iverson and Skinner]	140
Annual report for 1948 by State	804
Quarterly reports by State226,	
Frichinosis cases:	
Annual report for 1948 by State.	804
Quarterly reports by State	654

Tubercle bacilli, demonstration in sputum smears. Selection of choice particles compared with various sodium hydroxide concentration techniques	Page
[Spendlove, Cummings and Patnode]	574
monary calcifications [Beadenkopf, Loosli, Lack, Rice and Slattery] Tuberculosis:	17
Cases: Annual report for 1948 by State	004
	804
Quarterly reports by State 226	
Control in the Philippines [Young].	123
Control program of nine California mental institutions [Oechsli]	4
Deaths—White and nonwhite	708
Diagnosis	574
Mental hospitals—editorial [Anderson]	1
Mortality in older age groups—editorial [Anderson]	403
Mortality in the United States, 1947 [Lewis]	405
Mortality rates associated with selected characteristics of patient	
population	687
Nursing in—editorial [Anderson]	121
Patients discharged from sanatoria, studies	687
Pneumonectomy followed by immediate thoracoplasty [Iverson and	
Skinner]	140
Research in Muscogee County, Ga 259	, 263
Social problems	685
Studies in Muscogee County, Ga:	
I. Community-wide tuberculosis research [Comstock]	259
II. X-ray findings in a community-wide survey and its coverage	
as determined by a population census [Burke, Schenck and	
Thrash]	263
Thoracoplasty following pneumonectomy [Iverson and Skinner]	140
Tularemia cases:	
Annual report for 1948 by State	804
Quarterly reports by State226	, 654
Weekly State reports	34,
64, 88, 112, 154, 195, 218, 246, 296, 325, 349, 392, 434, 462, 518, 540, 584, 616, 644, 676, 710, 742, 778,	
Typhoid and paratyphoid fever:	
Cases:	
Annual report for 1948 by State	804
Quarterly reports by State226	
Weekly State reports	34,
64, 88, 112, 154, 195, 218, 246, 296, 325, 349, 392, 434, 462,	
518, 540, 584, 616, 644, 676, 710, 742, 778,	
Colombia	436
Typhus fever:	
Endemic, cases:	
Annual report for 1948 by State	804
Quarterly reports by State226,	
Foreign reports	-38,
68, 92, 118, 200, 256, 302, 354, 398, 469, 498, 522, 548, 587, 683, 716, 747, 784,	650,
Murine, in Louisville, Ky. [Good and Kotcher]	229

U

	· ·	
Undulant fever cases:		Page
Annual report for 1948 by St		
Quarterly reports by State		226, 654
	V	
Venezuela: Influenza		
Vincent's infection cases:		
Annual report for 1948 by Sta	ste	804
Quarterly reports by State		226 654
Virgin Islands: Notifiable diseases		
Virus, poliomyelitis, flocculation to		
	w	
Whooping cough cases:	**	
Annual report for 1948 by Sta	ite	804
Quarterly reports by State		
Weekly State reports		34.
	95. 218. 246. 296. 3	25, 349, 392, 434, 462, 494,
		644, 676, 710, 742, 778, 808
World Health Organization influen		
	X	
X-ray screens and films, characte	ristics of commerci	al—IV, V, VI [Van
Allen]		430, 581, 706
	Y	
Yaws cases:		
Annual report for 1948 by Sta	te	806
Quarterly reports by State		228, 656
Yellow fever:		
Cases:		
Annual report for 1948 by	State	806
Quarterly reports by State	0	656
Foreign reports		68,
		54, 400, 469, 498, 522, 549, 20, 684, 716, 748, 784, 815
	001, 0	20, 002, 110, 140, 104, 810

Public Health Service Author Index

Includes Papers by Public Health Service Personnel Printed in Public Health Reports and Other Publications January-June 1949*

A

- Adamik, R. E.: See Endicott, K. M. (Gillman, Brecher, Ness, Clarke, and Adamik).
- Algire, G. H.: Blood pressure measurements and changes in peripheral vascular bed of unanesthetized mice (abstract). Federation Proc. 8: 349.
- ——— (and Legallais): A method for correlating blood pressure measurements with changes in the peripheral vascular bed of unanesthetized mice (abstract). Anat. Rec. 103: 566.
- ——— (Merwin and Legallais): Microscopic studies on tissue growth and vascular reactions in transparent chambers in mice (abstract). Anat. Rec. 103: 590.
- Altenderfer, Marion E. (and Crowther): Relationship between infant mortality and socioeconomic factors in urban areas. Pub. Health Rep. 64: 331.
- Altland, Paul D.: Effect of discontinuous exposure to 25,000 feet simulated altitude on growth and reproduction of the albino rat. J. Exper. Zool. 110: 1.
- Effect of rest intervals on maintenance of previously induced high altitude acclimatization (abstract). Federation Proc. 8: 3.
- Anderson, Evelyn (and Haymaker): Carbohydrate metabolism in the decerebrated rat (abstract). Federation Proc. 8: 4.
- ———— Glucose tolerance in decerebrated rats after relatively long survival. Proc. Soc. Exper. Biol. and Med. 70: 86.
- Anderson, Robert J.: Interrelation of social problems. Editorial. Pub. Health Rep. 64: 685.
 - Nursing in tuberculosis. Editorial. Pub. Health Rep. 64: 121.
 - Tuberculosis in mental hospitals. Editorial. Pub. Health Rep. 64: 1.
- Tuberculosis mortality in older age groups. Editorial. Pub. Health Rep. 64: 403.
- Andrews, Howard L.: Instruments in radiobiology. Nucleonics 4: 109.
- Radiation hazards and industrial hygiene. Indust. Hyg. Newsletter
 9: 4, July.

NOTE: Address requests for reprints from Public Health Reports, Journal of Venereal Disease Information, and the Journal of the National Cancer Institute to the Surgeon General, Public Health Service, Washington 25, D. C.

^{*}All items marked with an asterisk were published in 1948, but reported too late for inclusion in Index to vol. 63, part I.

*Apter, Nathaniel S. (Eisele, McCullough, and Halstead): Cerebral pathology in brucellosis. A combined medical, neuropsychiatric and experimental study. In Transactions of the American Neurological Association, Richmond, Va., William Byrd Press, p. 39.

Arnold, F. A., Jr.: Fluorine: Systemic, oral environmental, experimental, and control aspects. In Dentistry in public health by W. J. Pelton and J. M.

Wisan, Philadelphia, W. B. Saunders Co., p. 163.

—— See also Dean, H. Trendley (and Arnold); McCauley, H. Berton (Likins and Arnold); McCleod, Charlotte (and Arnold).

Arnold, R. C. (Mahoney, Nicholson, and Wright): Penicillin therapy of early syphilis: IV. J. Ven. Dis. Inform. 30: 125.

* (and Nicholson): Treatment failures versus reinfections following penicillin therapy in early syphilis. In Recent advances in the study of venereal diseases. A symposium, Raleigh, N. C., Venereal Disease Education Institute in cooperation with the United States Public Health Service, p. 107.

Ashburn, L. L. (Baker and Faulkner): Failure of thyroidectomy and thiouracil to protect rat liver from acute carbon tetrachloride injury. Proc. Soc. Exper.

Biol. and Med. 70: 624.

Aykroyd, W. R. (Jolliffe, Lowry, Moore, Sebrell, Shank, Tisdall, Wilder, and Zameenik): Medical resurvey of nutrition in Newfoundland, 1948. Canadian Med. Assoc. J. 60: 329.

B

Baker, W. H.: See Ashburn, L. L. (Baker and Faulkner).

Bartgis, Ida Louise: See Reardon, Lucy V. (and Bartgis).

Bateman, Jeanne C. (Shorr and Elgvin): Hypervolemic anemia in cirrhosis. J. Clin. Invest. 28: 539.

—— See also Rodriquez, Jack (Schwemlein, Bauer, Plotke, Peters, Kendell, and Rodriquez).

Beck, M. Dorothy (Bell, Shaw, and Huebner): Q fever studies in southern California. II. An epidemiological study of 300 cases. Pub. Health Rep. 64: 41.

—— See also Huebner, Robert J. (Jellison and Beck).

Bell, E. J. (Parker and Stoenner): Q fever. Experimental Q fever in cattle. Am. J. Pub. Health 39: 478.

---- See also Parker, Ralph R. (Bell and Stoenner).

Bell, J. Frederick (Wright and Habel): Rabies vaccine freed of the factor causing allergic encephalitis. Proc. Soc. Exper. Biol. and Med. 70: 457.

——— See also Habel, Karl (Bell and Wright); Habel, Karl (Endicott, Bell, and Spear).

Bell, Joseph A. (Pittman and Olson): Pertussis and aureomycin. Pub. Health Rep. 64: 589.

See also Beck, M. Dorothy (Bell, Shaw, and Huebner).

Benjamin, Edward A. (and Roberson): Some socioeconomic aspects of venereal disease among Negroes. Nat. Negro Health News 17: 11, Jan.-June.

Bennett, Harry J.: See Newton, Walter L. (Bennett and Figgat).

Berdan, Elsie T.: Better use of nursing personnel. Mod. Hosp. 72: 80.

Binford, Chapman H. (and Spriggs): Clinicopathologic conference. Am. J. Clin. Path. 19: 388.

—— See also Gledhill, Emerson Y. (Spriggs and Binford). 890594—50——3 Birren, James E.: See Fox, Charlotte (and Birren).

*Blanco, F. Leon (and Fite): The effect of fixatives on straining procedures for lepra bacilli in tissues. Internat. J. Leprosy 16: 367.

*Block, Louis: There is no comparison of costs without uniform accounting procedure. Mod. Hosp. 71: 79.

Block, Matthew: The development of intra-ocular transplants of embryonic rat liver (abstract). Anat. Rec. 103: 425.

Bloomfield, J. J.: Industrial hygiene highlights at APHA Meeting, Boston. Indust. Hyg. Newsletter 9: 3, Jan.

——— Today's legacy, tomorrow's signpost. Indust. Hyg. Newsletter 9: 10, June.

Blount, H. C. (and Smith): The influence of thyroid and thiouracil on mice exposed to roentgen radiation. Science 109: 83.

Board, Leonard M.: The home environment and accidents. Pub. Health Rep. 64: 383.

Bond, Howard W.: See Hundley, James M. (and Bond).

Boone, Bert R. (Randak, Ellinger, and Oppenheimer): Electrokymographic studies of the ventricular isometric relaxation phase of the cardiac cycle in man. J. App. Physiol. 1: 534:

Boone, Howard: See Gray, A. L. (Boone and Hibbets).

Borbiro, M. (and Szent, Györgyi): On the relation of tension and ATP in cross-striated muscle. Biol. Bull. 96: 162.

Bossak, Hilfred N.: An apparatus for the aseptic collection and dispensing of sheep blood. Am. J. Clin. Path. 19: 496.

——— (Rosenberg and Harris): A quantitative turbidimetric method for the determination of spinal fluid protein. J. Ven. Dis. Inform. 30: 100.

Bowdoin, C. D. (Henderson, Davis, Remein, and Morse): Socioeconomic factors in syphilis, Savannah, Ga. J. Ven. Dis. Inform. 30: 131.

Bowen, William J.: The absorption spectra and extinction coefficients of myoglobin. J. Biol. Chem 179: 235.

—— A method for analyzing dog muscles for mycglobin (abstract). Federation Proc. 8: 14.

Bowman, Richard W.: See Cannefax, George R. (Parkhurst and Bowman).

*Bradley, G. H.: Mosquito control in the United States. In Proc. 4th Internat. Cong. Trop. Med. and Malaria, Washington, D. C., U. S. Government Printing Office, vol. 2, p. 1598.

Brady, Frederick J.: International cooperation against the tsetse. Department of State Bull. 20: 722.

Brecher, G. (and Wexler): Obesity in albino mice due to single injections of gold-thioglucose. Proc. Soc. Exper. Biol. and Med. 70: 498.

—— See also Endicott, K. M. (Gillman, Brecher, Ness, Clarke, and Adamik).
Brewster, Agnes W. (and Fletcher): Studies of patients discharged from tuber-culosis sanatoria. II. Mortality rates associated with selected characteristics of the patient population. Pub. Health Rep. 64: 687.

Brinton, Hugh P.: See Fairhall, Lawrence T. (Keenan and Brinton). Brodsky, William A.: See Rapoport, S. (Brodsky, West, and Mackler).

Bucca, Matthew A.: See Carpenter, Charles M. (Bucca, Buck, Casman, Christensen, Crowe, Drew, Hill, Lankford, Morton, Peizer, Shaw, and Thayer).

Burch, Thomas A.: Experimental therapy of onchocerciasis with suramin and hetrazan. Bol. Ofic. san. panam. 28: 233.

Burgess, Robert W.: See Cooper, W. Clark (Ruhe, Coatney, Josephson, Young, and Burgess).

Burgoyne, F. H. (Heston, Hartwell, and Stewart): Cutaneous melanin production in the mouse by 5,9,10-trimethyl, 1-2-benzanthracene (abstract). Federation Proc. 8: 351.

Burk, Dean: See DuBuy, H. G. (Woods, Burk, and Lackey); Fiala, Silvio (and Burk); Hearon, John Z. (Burk and Schade); Riley, Vernon T. (Hesselbach, Fiala, Woods, and Burk); Wight, Kent (and Burk); Woods, Mark W. (Du-Buy, Burk, and Hesselbach).

Cannefax, George R.: A stable control serum for standardizing the sensitivity of tests used in the diagnosis and control of syphilis. J. Ven. Dis. Inform. 30: 169.

- (Parkhurst and Bowman): Comparative studies on mailed spinal fluid specimens, J. Ven. Dis. Inform. 30: 11.

Carpenter, Charles M.: (Bucca, Buck, Casman, Christensen, Crowe, Drew, Hill, Lankford, Morton, Peizer, Shaw, and Thayer): Evaluation of twelve media for the isolation of the gonococcus. Am. J. Syph., Gonor., and Ven. Dis. 33: 164.

Caster, W. O. (and Mickelsen): Variability in the Beckman spectrophotometer (abstract). Federation Proc. 8: 190.

Chalkley, Harold W.: A method for the measurement of average nuclear volumes independent of shape (abstract). Anat. Rec. 103: 433.

Chang, Yao-teh: See Smith, Milton I. (Jackson, Chang, and Longnecker).

Chapman, A. L.: The public health worker in conservation of vision for old age. Sight-Saving Rev. 19: 3.

Chapman, Kenneth W.: See Vogel, Victor H. (Chapman and Isbell).

Clark, Ariel M.: See Spicer, Samuel S. (Hanna and Clark); Spicer, Samuel S. (Wooley and Clark).

Clark, R. J. (and Firminger): Coarctation of the aorta associated with Adams-Stokes syndrome, complete heart block and bicuspid calcareous aortic valve. New England J. Med. 240: 710.

Clarke, F. A.: See Endicott, K. M. (Gillman, Brecher, Ness, Clarke, and Adamik). Coatney, G. Robert: Chemotherapy of malaria. Bol. Ofic. san. panam. 28: 27.

(and Cooper): Recrudescence and relapse in vivax malaria. In Proc. 4th Internat. Cong. Trop. Med. and Malaria, Washington, D. C., U. S. Government Printing Office, vol. 1, p. 629.

(Ruhe, Cooper, Josephson, and Young): Studies in human malaria. X. The protective and therapeutic action of chloroquine (SN 7618) against

St. Elizabeth strain vivax malaria. Am. J. Hyg. 49: 49.

See also Cooper, W. Clark (Ruhe, Coatney, and Josephson); Cooper, W. Clark (Ruhe, Coatney, Josephson, and Young); Cooper, W. Clark (Ruhe, Coatney, Josephson, Young, and Burgess); Rein, Charles R. (Bukantz, Kent, Cooper, Ruhe, and Coatney); Ruhe, David S. (Cooper, Coatney, and Josephson); Ruhe, David S. (Cooper, Coatney, Josephson, and Young).

Cohn, Jerome E. (and Shock): Blood volume studies in middle-aged and elderly

Am. J. Med. Sc. 217: 388.

Cole, H. N. (Plotke, Thomas, and Jenkins): Penicillin treatment of syphilis in pregnancy. J. Ven. Dis. Inform. 30: 95.

Coleman, Nell: See Haas, Victor H. (Wilcox and Coleman).

Cooper, W. Clark: Summary of antimalarial drugs. Pub. Health Rep. 64: 717. (Ruhe, Coatney, and Josephson): Studies in human malaria. XIII. The therapeutic action of metachloridine (SN 11, 437) against St. Elizabeth strain vivax malaria. Am. J. Hyg. 49: 355.

Cooper, W. Clark (Ruhe, Coatney, Josephson, and Young): Studies in human malaria. VIII. The protective and therapeutic action of quinacrine against St. Elizabeth strain vivax malaria. Am. J. Hyg. 49: 25.

(Ruhe, Coatney, Josephson, Young, and Burgess): Studies in human malaria. XI. The protective and therapeutic action of SN 6771 against St.

Elizabeth strain vivax malaria. Am. J. Hyg. 49: 60.

—— See also Coatney, G. Robert (and Cooper); Coatney, G. Robert (Ruhe, Cooper, Josephson, and Young); Rein, Charles R. (Bukantz, Kent, Cooper, Ruhe, and Coatney); Ruhe, David S. (Cooper, Coatney, and Josephson); Ruhe, David S. (Cooper, Coatney, Josephson, and Young).

*Corcoran, Mary E.: The psychological basis of personal problems of workers.

Indust. Nursing 7: 16.

Cram, Eloise B.: See Wright, Willard H. (and Cram).

Crowther, Beatrice: See Altenderfer, Marion E. (and Crowther).

Culbertson, James T.: Plans for United States cooperation with the World Health Organization in the international influenza study program. Am. J. Pub. Health 39: 37.

Cummings, Martin M.: The laboratory diagnosis of tuberculosis. Am. J. Pub. Health 35: 361.

- See also Spendlove, George A. (Cummings and Patnode).

D

Dalton, Albert J. (Kahler, Lloyd, and Striebich): A smear technique for preparing cells for study under the electron microscope (abstract). Anat. Rec. 103: 437.

— (Kahler, Kelly, Lloyd, and Striebich): Some observations on the mitochondria of normal and neoplastic cells with the electron microscope. J. Nat. Cancer Inst. 9: 439.

Davies, D. F.: Age and osmotic work in urea excretion (abstract). Federation Proc. 8: 32.

Davis, Bernard D.: Isolation of biochemically deficient mutants of bacteria by means of penicillin (abstract). Federation Proc. 8: 194.

Davis, Burnet M.: The British National Health Service. Pub. Health Rep. 64: 161.

*Davis, Dorland J.: Complement fixation reaction in psittacosis. Arch. Int. Med. 81: 623.

——— (and Pittman): Effectiveness of streptomycin in treatment of experimental conjunctivitis caused by *Hemophilus sp.* Am. J. Ophth. 32: 111, June, pt. 2.

——— (and Vogel): Recovery of psittacosis virus from chicks hatched from inoculated eggs. Proc. Soc. Exper. Biol. and Med. 70: 585.

Davis, James O. (and Shock): Theophylline ethylene diamine and renal function in control subjects and decompensated cardiac patients (abstract). Federation Proc. 8: 32.

Davis, Warren T., Jr.: See Bowdoin, C. D. (Henderson, Davis, Remein, and Morse).

Dean, H. Trendley: Fluorine and dental caries. Illinois Med. J. 95: 33.

----- Fluorine in dental caries control. Postgrad. Med. 5: 361.

Fluorine: Water-borne fluorides and dental health. In Dentistry in public health by W. J. Pelton and J. M. Wisan, Philadelphia, W. B. Saunders Co., p. 136.

Dean H. Trendley (and Arnold): Dental research and the National Institutes of Health: I. Intramural research. J. Am. Dent. Assoc. 38: 15.

Deibert, Austin V.: Cancer control: A comprehensive approach. Hospitals 23: 49. May.

The dentist in the national program of cancer control. Am. J. Pub. Health 39: 772.

For early cancer recognition: New films to dramatize control. Hospitals 23: 66, Feb.

——— Professional education for cancer control. Pub. Health Rep. 64: 749.

Dent, Jack E. (Morlan and Hill): Effects of DDT dusting on domestic rats under colony and field conditions. Pub. Health Rep. 64: 666.

Dhyse, Frederick G.: See Hertz, Roy (Dhyse and Tullner).

Diamond, Murray: See McDaniel, T. W. (and Diamond).

Dormin, K. Barbara: Public health nurse in diabetes control. Pub. Health Nursing 41: 9.

Dorn, Harold F.: Forms that meet current needs. How a standardization committee simplified, in format and number, the clinical records for Federal hospitals. Hospitals 23: 62, Jan.

- Morbidity from cancer. J. Washington Acad. Sc. 39: 117.

The use of statistics in cancer control programs. Am. J. Pub. Health 39: 602.

Dow, Richard P.: See Maier, Paul P. (and Dow).

*Drake, Roy A. (and Geil): Homosexual activity; Relation of degree and role to the Goodenough test and to the Cornell Selectee Index. J. Nerv. and Ment. Dis. 108: 217.

Dreessen, W. C.: Pneumoconiosis in the United States. In Proc. 9th Internat. Cong. Indust. Med., Bristol, England, John Wright and Sons Ltd., p. 720.

DuBuy, H. G. (Woods, Burk, and Lackey): Enzymatic activities of isolated amelanotic and melanotic granules of mouse melanomas and a suggested relationship to mitochondria. J. Nat. Cancer Inst. 9: 325.

- See also Woods, M. W. (DuBuy, Burk, and Hesselbach).

Dunn, Thelma B.: Blood dyscrasia in mice ingesting toluidine blue (abstract). Federation Proc. 8: 353.

——— Some observations on the normal and pathologic anatomy of the kidney of the mouse. J. Nat. Cancer Inst. 9: 285.

Dyer, R. E.: Q fever. History and present status. Am. J. Pub. Health 39: 471.

E

Eagle, Harry: Host, drug, and parasitic factors that modify the therapeutic activity of penicillin. In Advances in internal medicine, New York City, Interscience Press, vol. 3, p. 105.

(and Fleischman): The antibody response in rabbits to killed suspensions of pathogenic T. pallidum. In Recent advances in the study of venereal diseases. A symposium, Raleigh, N. C., Venereal Disease Education Institute

in cooperation with the United States Public Health Service, p. 17.

* Therapeutic activity of bacitracin in rabbit syphilis, and its synergistic action with penicillin. A preliminary report. In Recent advances in the study of venereal diseases. A symposium, Raleigh, N. C., Venereal Disease Education Institute in cooperation with the United States Public Health Service, p. 320.

- Eagle, Harry (Fleischman and Musselman): The serum concentration of penicillin G in mice, rabbits, and men after its intramuscular injection in aqueous solution. J. Bact. 57: 119.
- Earle, W. R. (Evans, Sanford, Likely, and Schilling): Certain recently developed tissue culture procedures utilizing perforated cellophane as a culture substrate (abstract). Anat. Rec. 103: 443.
- Eddy, Nathan B.: Cooperation in narcotics. Drug and Allied Industries 5: 8.

 —— Metopon hydrochloride. An experiment in clinical evaluation. Pub.

 Health Rep. 64: 93.
- * Progress in drug therapy of pain. Am. Prof. Pharmacist 14: 252.
- Eisenberg, Henry (and Plotke): Penicillin in oil and beeswax in the treatment of syphilis in clinic patients. Am. J. Med. 6: 449.
- (Plotke and Baker): Asexual syphilis in children. J. Ven. Dis. Inform. 30: 7.
- Elias, Hans: The liver. I. Normal histology. II. Histopathology of portal cirrhoses (abstract). Anat. Rec. 103: 591.
- The network of bile capillaries and its relation to the hepatic parenchyma (abstract). Anat. Rec. 103: 444.
- Ellinger, George F.: See Boone, Bert R. (Randak, Ellinger, and Oppenheimer). Emmart, Emily W. (Kissling and Stark). The tuberculostatic action of nocardia extracts in white mice. J. Bact. 57: 509.
- —— (McClosky, Smith, and Lieberman): Acquired resistance of M. tuberculosis to streptomycin in vitro and its behavior in experimental infections. Am. Rev. Tuberc. 59: 438.
- See also Smith, M. I. (and Emmart).
- *Emmons, Chester W.: Coccidoidomycosis and haplomycosis. In Proc. 4th Internat. Cong. Trop. Med. and Malaria, Washington, D. C., U. S. Government Printing Office, vol. 2, p. 1278.
- Diagnostic problems in medical mycology. Am. J. Pub. Health 39: 713.
 Histoplasmosis in animals. Tr. New York Acad. Sc. 11: 248.
- Endicott, K. M. (Gillman, Brecher, Ness, Clarke, and Adamik): A study of
- Erickson, Paul T.: La estreptomicina como auxiliar terepéutico de las sulfonas en ciertas manifestaciones leprosas (Streptomycin. A useful adjunct to sulfones in certain lesions). America Clinica 14: 123.
- Errara, Maurice: Liver glutaminases. J. Biol. Chem. 178: 483.
- —— (and Greenstein): Phosphate-activated glutaminase in kidney and other tissues. J. Biol. Chem. 178: 495.
- Eschenbrenner, Allen: Injury and recovery of mouse testes following whole-body exposure to X-radiation (abstract). Federation Proc. 8: 354.
- Evans, Virginia J.: See Earle, W. R. (Evans, Sanford, Likely, and Schilling).
- Ewing, William H.: The relationship of bacterium antiratum and members of the tribe Mimeae (DeBord). J. Bact. 57: 659.
- Shigella nomenclature. J. Bact. 57: 633.

F

- Fairhall, Lawrence T.: Industrial toxicology. Mercury. Indust. Hyg. Newsletter 9: 10, Mar.
- --- Industrial toxicology. Phosgene. Indust. Hyg. Newsletter 9: 13, Jan.
- ——— (Keenan and Brinton): Cobalt and the dust environment of the cemented tungsten carbide industry. Pub. Health Rep. 64: 484.

Fales, W. Thurber (and Moriyama): International adoption of principles of morbidity and mortality classification. Am. J. Pub. Health 39: 31.

Farber, Herbert R. (Yiengst and Shock): The effect of therapeutic doses of aspirin on the acid-base balance of the blood in normal adults. Am. J. Med. Sc. 217: 256.

*Farmer, Thomas W.: The Jarisch-Herxheimer reaction in early syphilis treated with crystalline penicillin G. In Recent advances in the study of venereal diseases. A symposium, Raleigh, N. C., Venereal Disease Education Institute in cooperation with the United States Public Health Service, p. 121.

———— (and Ravitch): The fate of Wassermann antibody passively transferred by intravenous injection of plasma into human subjects. *In* Recent advances in the study of venereal diseases. A symposium, Raleigh, N. C., Venereal Disease Education Institute in cooperation with the United States Public Health Service, p. 329.

——— See also Hoekenga, Mark T. (and Farmer); Pardo, Oswaldo A. (and Farmer); Tucker, Harold A. (Farmer and Fein).

Faulkner, R. R.: See Ashburn, L. L. (Baker and Faulkner).

*Felix, Robert H.: A state-wide mental health program. In Proc. Sixty-second Annual Meeting, Conference of State and Provincial Health Authorities of North America, Chicago, The Conference, p. 25.

The teacher's role in mental health. School Life 31: 1.

Felton, Lloyd D.: The significance of antigen in animal tissues. J. Immunol. 61: 107.

—— (Prescott, Kauffmann, and Ottinger): Polysaccharides isolated from tissues of normal, immune, and immunologically paralyzed mice (abstract). Federation Proc. 8: 402.

A possible conjugated [polysaccharide-protein] antigen of the Diplococcus pneumoniae (abstract). In Proc. 49th Meeting, Soc. Am. Bact., p. 88.

Ferguson, Charles: Hypospadias: A method to improve the Chordee effect. J. Urol. 61: 1052.

Fiala, Silvio: Glycolysis and the Pasteur effect on the subcellular level (abstract). Federation Proc. 8: 197.

—— (and Burk): On the mode of iron binding by siderophilin, conalbumin, hydroxylamine, aspergillic acid, and other hydroximic acids. Arch. Biochem. 20: 172.

- See also Riley, Vernon T. (Hesselbach, Fiala, Woods, and Burk).

Figgat, William B.: See Newton, Walter L. (Bennett and Figgat).

Firminger, H. I.: See Clark, R. J. (and Firminger); Mulay, A. S. (and Firminger). Fite, G. L.: See Blanco, F. Leon (and Fite).

Fitts, Charles N.: The National Institutes of Health. Nat. Negro Health News 17: 1, Jan.-June.

Fletcher, Ralph Carr: See Brewster, Agnes W. (and Fletcher).

Fodor, Paul J. (Price and Greenstein): Preparation of L-, and D- alanine by enzymatic resolution of the acetyl-DL-alanine. J. Biol. Chem. 178: 503.

Fones, William S. (and White): Preparation of p-dimethylaminoazobenzene containing isotopic nitrogen. Arch. Biochem. 20: 118.

Ford, Malcolm J.: See Wilkerson, Hugh L. C. (and Ford).

Forsbeck, Filip C.: "Infection unit" and "Index of aggregation." Suggested epidemiological terms. Pub. Health Rep. 64: 343.

Forsyth, Bruce D.: Importance of public health dentistry. Bol. Ofic. san. panam. 28: 56.

Forsyth, Bruce D.: The role of dentistry in the field of public health. Pennsylvania Dent. J. 16: 105.

Your own oral health. Colhecon 4: 8.

Fox, Charlotte (and Birren): Some factors affecting vocabulary size in later maturity: Age, education, and length of institutionalization. J. Gerontol. 4: 19.

Francis, Edward: Twenty-five year survival of a *Pasteurella pestis* culture without transfer. Pub. Health Rep. **64**: 238.

Frank, Karl: See Wikler, Abraham (and Frank).

*Frey-Wyssling, A. (Muhlethaler and Wyckoff): Mikrofibrillenbau der pflanzlichen Zellwände (Microfibrillary structure of cellular walls of plants). Experientia 4: 475.

*Friedenwald, Jonas S. (Buschke, Crowell, and Hollaender): Effects of ultraviolet irradiation on the corneal epithelium. II. Exposure to monochromatic radiation. J. Cell. and Comp. Physiol. 32: 161.

Friedkin, Morris (and Lehninger): Oxidation-coupled incorporation of inorganic radiophosphate into phospholipide and nucleic acid in a cell-free system. J. Biol. Chem. 177: 775.

Furculow, Michael L. (and Ruhe): Histoplasmin sensitivity among cattle. Am. J. Pub. Health 39: 719.

G

Gafafer, William M.: Sickness absenteeism among industrial workers. Third and fourth quarters of 1948. Pub. Health Rep. 64: 641.

*Geil, George A.: The Goodenough test as applied to adult delinquents. J. Clin. Psychopath. 9: 62.

See also Drake, Roy A. (and Geil).

Gerloff, R. K.: See Lackman, David B. (Parker and Gerloff).

Giacomo, Rosalie: See Trautman, J. A. (and Giacomo).

Gledhill, Emerson Y. (Spriggs and Binford): Needle aspiration in the diagnosis of lung carcinoma—Report of experience with seventy-five aspirations. Am. J. Path. 19: 235.

Goldstein, Marcus S.: Medical group practice in the United States: V. Growth of groups. Journal-Lancet 69: 42.

Good, Newell E. (and Kotcher): Murine typhus fever in Louisville, Kentucky. Pub. Health Rep. 64: 229.

Gordon, Donald A.: See Oliphant, John W. (Gordon, Meis, and Parker).

Gottschalk, Louis: See Grinker, Roy R. (and Gottschalk).

Gover, Mary: Statistical studies of heart disease. III. Heart disease associated with other major causes of death as primary or contributory cause. IV. Mortality from heart disease (all forms) related to geographic section and size of city. Pub. Health Rep. 64: 104; 439.

Gratch, I. (Purlia and Martin): Effect of sodium fluoroacetate (1080) in poisoned rats on plague diagnosis procedures. Preliminary report. Pub. Health Rep.

Gravatt, Margaret: See Halbert, Seymour P. (and Gravatt).

Gray, A. L. (Boone and Hibbets): VD case finding in Quitman County, Mississippi. J. Ven. Dis. Inform. 30: 127.

——— (Iskrant and Hibbets): Syphilis contact investigation in a rural county in Mississippi. J. Ven. Dis. Inform. 30: 165.

Greco, Antoinette: See Maver, Mary E. (and Greco).

Greco, J.: See Lillie, R. D. (Greco and Laskey).

Greenberg, Joseph: Hypersensitivity to sulfadiazine of a chlorguanide-resistant strain of *Plasmodium gallinaceum*. J. Nat. Malaria Soc. 8: 80.

Inhibition of the antimalarial activity of chlorguanide by pteroylglutamic acid. Proc. Soc. Exper. Biol. and Med. 71: 306.

—— See also Josephson, Edward S. (Taylor, Greenberg, and Nadel); Nadel, Eli M. (Taylor, Greenberg, and Josephson).

Greenstein, Jesse P. (and Leuthardt): Dehydropeptidase activity in tumors. J. Nat. Cancer Inst. 9: 389.

——— (and Price): α-Keto acid-activated glutaminase and asparaginase. J. Biol. Chem. 178: 695.

——— See also Errara, Maurice (and Greenstein); Fodor, Paul J. (Price and Greenstein); Price, Vincent E. (Meister and Greenstein).

*Griffith, Robert L.: Chylous ascites and chylothorax—Report of a case in which the cause was undetermined. Northwest Med. 47: 800.

Grinker, Roy R. (and Gottschalk): Headaches and muscular pains. Psychosomatic Med. 11: 45.

Guthrie, Robert: Studies on a purine-requiring mutant strain of Escherichia coli.
J. Bact. 57: 39.

H

Haas, Victor H. (Wilcox and Coleman): Modification of *Plasmodium gallinaceum* infections by certain tissue extracts. J. Nat. Malaria Soc. 8: 85.

Habel, Karl (Bell and Wright): Benzene-inactivated rabies vaccine. Proc. Soc. Exper. Biol. and Med. 70: 455.

——— (Endicott, Bell, and Spear): Immunological evidence on the role of the lymphocyte in antibody formation. J. Immunol. 61: 131.

See also Bell, J. Frederick (Wright and Habel).

Hagan, Thomas L. (and Cook): Dental health in Louisiana. J. Am. Dent. Assoc. 38: 31.

Halbert, Seymour P. (and Gravatt): Prevalence of antibiotic-producing coliform organisms. Further studies. Pub. Health Rep. 64: 313.

Halpin, Evelyn: Accident facts from accident fatalities. Pub. Health Rep. 64: 388.

Hanna, Charles H.: See Spicer, Samuel S. (Hanna and Clark).

Hare, W. V. (and Stewart): Spontaneous adenomatous gastritis of mice of strain dba (abstract). Federation Proc. 8: 357.

Harper, T. C.: See Michele, Arthur A. (and Harper).

Harris, Ad: See Bossak, Hilfred N. (Rosenberg and Harris).

Harris, M. R.: See von Oettingen, W. F. (Donahue, Yagoda, Monaco, and Harris).

Hartwell, J. L.: See Burgoyne, F. H. (Heston, Hartwell, and Stewart).

Hayes, Wayland J.: See Simmons, Samuel W. (and Hayes).

Hearon, John Z. (Burk and Schade): Physicochemical studies of reversible and irreversible complexes of cobalt, histidine, and molecular oxygen. J. Nat. Cancer Inst. 9: 337.

- See also Tschirgi, R. D. (Gerard, Jenerick, Boyarsky, and Hearon).

Heller, John R., Jr.: Cancer control: Mid-channel in cancer research. Hospitals 23: 48, May.

*—— The essentials of a State cancer program. In Proc. Sixty-second Annual Meeting, Conference of State and Provincial Health Authorities of North America. Chicago, The Conference, p. 50.

----- See also Hullerman, Hugo V. (and Heller).

- Henderson, Adele: The nurse in a mental health clinic. Pub. Health Nursing 41: 42.
- Heppel, Leon A.: Cytochrome C reductase (DPN) (abstract). Federation Proc. 8: 205.
- ———— (and Porterfield): Metabolism of inorganic nitrite and nitrate esters. I.

 The coupled oxidation of nitrite by peroxide-forming systems and catalase.

 J. Biol. Chem. 178: 549.
- See also Horecker, B. L. (and Heppel).
- Hertz, Roy (Dhyse and Tullner): The elevation of plasma riboflavin in estrogentreated female chicks. Endocrinology 44: 283.
- ——— (and Tullner): Inhibition of estrogen-induced growth in the genital tract of the female chick by a purine antagonist; Reversal by adenine. Science 109: 539.
- ———— Quantitative interference with estrogen-induced tissue growth by folic acid antagonists. Endocrinology 44: 278.
- Hesselbach, Marie L.: See Riley, Vernon T. (Hesselbach, Fiala, Woods, and Burk); Woods, Mark W. (DuBuy, Burk, and Hesselbach).
- Heston, W. E.: See Burgoyne, F. H. (Heston, Hartwell, and Stewart).
- Hibbets, Richard S.: See Gray, A. L. (Boone and Hibbets); Gray, A. L. (Iskrant and Hibbets).
- Highman, Benjamin: See Spicer, Samuel S. (Highman and Monaco).
- Hill, Elmer L: See Dent, Jack E. (Morlan and Hill).
- Hill, Justina: See Carpenter, Charles M. (Bucca, Buck, Casman, Christensen, Crowe, Drew, Hill, Lankford, Morton, Peizer, Shaw, and Thayer).
- Hingson, Robert A.: The development of the hypospray for parenteral therapy by jet injection. Anesthesiology 10: 66.
- *[Hingson, Robert A.:] Therapeutic nerve block. Physicians Bull. 13: 5.
- —— See also Hughes, James G. (Lovejoy, Lynn, and Hingson); Lull, Clifford B. (and Hingson).
- *Hoekenga, Mark T. (and Farmer): The Jarisch-Herxheimer reaction in neurosyphilis. In Recent advances in the study of venereal diseases. A symposium, Raleigh, N. C., Venereal Disease Education Institute in cooperation with the United States Public Health Service, p. 130.
- Hoge, Vane M. (and Steinle): State legislation for minimum standards of hospital maintenance and operation. Pub. Health Rep. 64: 69.
- Hogeboom, George H.: Cytochemical studies of mammalian tissues. II. The distribution of diphosphopyridine neucleotide-cytrochrome C reductase in rat liver fractions. J. Biol. Chem. 177: 847.
- Hollaender, Alexander: See Friedenwald, Jonas S. (Buschke, Crowell, and Hollaender).
- Hollis, Mark D.: Stream pollution. In Proc. 47th Annual Conference, Surgeon General, Public Health Service, and Chief, Children's Bureau with State and Territorial Health Officers, State Mental Health Authorities, and State Hospital Survey and Construction Authorities, Washington, D. C., Public Health Service, p. 41. Processed.
- Hope, Malcolm C.: Licensure works for the hospital. Mod. Hosp. 72: 75.
- Horecker, B. L.: Cytochrome C reductase (TPN) from liver (abstract). Federation Proc. 8: 207.
- ——— (and Heppel): The reduction of cytochrome C by xanthine oxidase. J. Biol. Chem. 178: 683.
- Howell, Stacey F.: Purification, crystallization and properties of circuiin, an antibiotic from the bacillus circulans group (abstract). Federation Proc. 8: 208.

Hubbard, John P. (Bain and Pennell): School health services. A report from the American Academy of Pediatries study of child health services. Am. J. Pub. Health 39: 781.

Hudgens, J. A.: See Lawrence, R. L. (and Hudgens).

Hudson, Claude S.: See Stewart, Laura C. (Richtmyer and Hudson).

Huebner, Robert J. (Jellison and Beck): Q fever—A review of current knowledge. Ann. Int. Med. 30: 495.

—— (Jellison, Beck, and Wilcox): Q fever studies in southern California. III. Effects of pasteurization on survival of C. burneti in naturally infected milk. Pub. Health Rep. 64: 499.

- See also Beck, M. Dorothy (Bell, Shaw, and Huebner).

Hueper, Wilhelm C.: Environmental and occupational cancer. Pub. Health Rep. Supp. No. 209, 69 pp.

——— Prevention and treatment of atherosclerosis. In Medical clinics of North America (New York number), Philadelphia, W. B. Saunders Co., p. 773.

Problems in the control of occupational cancer. Indust. Med. 18: 163.
*Hughes, James G. (Lovejoy, Lynn, and Hingson): The use of continuous caudal analgesia for control of hypertension in acute nephritis. Am. J. Dis. Child. 75: 291.

*Hughes, John H.: Public Health Service foreign quarantine entomology relating to aircraft. Proc. Florida Anti-mosquito Assoc. 19: 70.

Hullerman, Hugo V. (and Heller): Organized attack on syphilis. A survey of hospital participation. Hospitals 23: 59, Feb.

Hundley, James M.: Influence of intestinal bacteria on synthesis of nicotinic acid from tryptophan. Proc. Soc. Exper. Biol. and Med. 70: 592.

——— Role of the gastrointestinal tract in the conversion of tryptophan to nicotinic acid (abstract). Federation Proc. 8: 386.

——— (and Bond): A study of the conversion of tryptophan to N'-Methylnicotinamide using isotopic carbon. Arch. Biochem. 21: 313.

Hunt, G. Halsey: Prepayment plans for medical care: History and present status. Med. Ann. District of Columbia 18: 311.

I

*Isbell, Harris: Methods and results of studying experimental human addiction to the newer synthetic analgesics. Ann. New York Acad. Sc. 51: 108.

*—— The newer analgesic drugs; Their use and abuse. Ann. Int. Med. 29: 1003.

——— (and Vogel): The addiction liability of methadon (amidone, dolophine, 10820) and its use in the treatment of the morphine abstinence syndrome. Am. J. Psychiat. 105: 909.

- See also Vogel, Victor H. (Chapman and Isbell).

Iskrant, Albert P.: See Bauer, Theodore J. (and Iskrant); Gray, A. L. (Iskrant and Hibbets); Spoto, Joseph F. (and Iskrant).

Iverson, Roland K. (and Skinner): Pneumonectomy followed by immediate thoracoplasty. Preliminary report of three cases. Pub. Health Rep. 64: 140.

J

Jackson, Ernest L.: See Smith, Milton I. (Jackson, Chang, and Longnecker).
Jacobson, Leon O. (Marks and Lorenz): Hematological effects of ionizing radiations.
Radiology 52: 371.

Jellison, William L.: See Huebner, Robert J. (Jellison and Beck); Huebner, Robert J. (Jellison, Beck, and Wilcox). Jenkins, Kenneth H.: See Cole, H. N. (Plotke, Thomas, and Jenkins).

Johnston, Dorothy: Educational program at the Alto Medical Center-Alto,

Georgia (Report). J. Ven. Dis. Inform. 30: 120.

Josephson, Edward S. (Taylor, Greenberg, and Nadel): The ascorbic acid content of the adrenal glands of chicks infected with Plasmodium gallinaceum. J. Nat. Malaria Soc. 8: 132.

- See also Coatney, G. Robert (Ruhe, Cooper, Josephson, and Young); Cooper, W. Clark (Ruhe, Coatney, and Josephson); Cooper, W. Clark (Ruhe, Coatney, Josephson, and Young); Cooper, W. Clark (Ruhe, Coatney, Josephson, Young, and Burgess); Nadel, Eli M. (Taylor, Greenberg, and Josephson); Ruhe, David S. (Cooper, Coatney, and Josephson); Ruhe, David S. (Cooper, Coatney, Josephson, and Young).

K

Kahler, Herbert: See Dalton, Albert J. (Kahler, Kelly, Lloyd, and Striebich); Dalton, Albert J. (Kahler, Lloyd, and Striebich); Lloyd, B. J. (and Kahler). Kaiser, Raymond F.: Cancer teaching in dental schools. Pub. Health Rep. 64:

Cancer teaching in medical schools. Pub. Health Rep. 64: 757.

Kaplan, Henry S. (and Murphy): The effect of local roentgen irradiation on the biological behavior of a transplantable mouse carcinoma. I. Increased frequency of pulmonary metastasis. J. Nat. Cancer Inst. 9: 407.

Kauffmann, Gladys: See Felton, Lloyd D. (Kauffmann and Ottinger); Felton, Lloyd D. (Prescott, Kauffmann, and Ottinger).

Keenan, Robert G.: See Fairhall, Lawrence T. (Keenan and Brinton).

Kelly, M. G.: See Dalton, Albert J. (Kahler, Kelly, Lloyd, and Striebich).

Kent, John F.: See Rein, Charles R. (Bukantz, Kent, Cooper, Ruhe, and Coatney). Kiefer, Norvin C.: Public health planning for national defense. In Proc. 47th Annual Conference, Surgeon General, and Chief, Children's Bureau with State and Territorial Health Officers, State Mental Health Authorities, and State Hospital Survey and Construction Authorities, Washington, D. C., Public Health Service, p. 48. Processed.

Kissling, R. E.: See Emmart, E. W. (Kissling and Stark).

Knutson, J. W.: Evaluation of the use of fluorine as a caries control measure; The effect of topically applied fluorides. J. Am. Dent. Assoc. 38: 204.

Kohls, Glen M.: See Philip, Cornelius B. (and Kohls).

*Komp, William H. W.: The anopheline vectors of malaria of the world. In Proc. 4th Internat. Cong. Trop. Med. and Malaria, Washington, D. C., U. S. Government Printing Office, vol. 1, p. 614.

Kornberg, A.: Inorganic pyrophosphate in the enzymatic synthesis of DPN (abstract). Federation Proc. 8: 215.

Krueger, F. J.: See Michele, Arthur A. (and Krueger).

L

Labaw, L. W.: See Smith, Falconer (and Labaw).

Lackey, Mary D.: See DuBuy, H. G. (Woods, Burk, and Lackey).

Lackman, David B. (Parker and Gerloff): Serological characteristics of the pathogenic rickettsia occurring in Amblyomma maculatum ticks (abstract). In Proc. 49th General Meeting Soc. Am. Bact., p. 84.

Laki, Koloman: Transition of fibrinogen to fibrin (abstract). Federation Proc.

Laskey, Alice M.: See Lillie, R. D. (Greco and Laskey).

Lawrence, R. L. (and Hudgens): Intestinal parasites in a prison population. Mil. Surgeon 104: 371.

Lear, Walter J.: The industrial hygienist and medical care. Indust. Hyg. Newsletter 9: 4, Jan.

Legallais, Frances Y.: See Algire, G. H. (and Legallais); Algire, G. H. (Mervin and Legallais).

Leuthardt, Florence M.: See Greenstein, Jesse P. (and Leuthardt).

Lewis, John A., Jr.: Problems in out-patient management of syphilis in the South. J. Ven. Dis. Inform. 30: 19.

Lewis, Sara A.: Tuberculosis mortality in the United States, 1947. Pub. Health Rep. 64: 405.

Lieberman, J. E.: See Emmart, Emily W. (McClosky, Smith, and Lieberman). Likely, Gwendolyn D.: See Earle, Wilton R. (Evans, Sanford, Likely, and Schilling).

Likins, R. C.: See McCauley, H. Berton (Likins and Arnold).

Liljegren, E. J.: See Webster, Stewart H. (Liljegren and Zimmer).

Lillie, R. D.: On the destruction of cytoplasmic basophilia (ribonucleic acid) and of the metrachromatic basophilia of cartilage by the glycogen splitting enzyme malt diastase: A histochemical study. Anat. Rec. 103: 611.

——— (Greco, and Laskey): Histochemical studies with malt diastase; Partial separation of the fractions destroying cytoplasmic basophilia (ribonucleic acid) and the metachromasia of cartilage. Anat. Rec. 103: 635.

*Lloyd, B. J., Jr. (and Kahler): Sedimentation of the chicken tumor I agent (abstract). Acta Unio Internationalis Contra Cancrum 6: 209.

—— See also Dalton, A. J. (Kahler, Kelly, Lloyd, and Striebich); Dalton, Albert J. (Kahler, Lloyd, and Striebich).

Long, Constance: The director of nursing service. Am. J. Nursing 49: 354.

*—— Nursing costs more than it should. Mod. Hosp. 71: 80.

Longnecker, William H.: See Smith, Milton I. (Jackson, Chang, and Longnecker). Lorenz, Egon: See Jacobson, Leon O. (Marks and Lorenz).

Lowry, James V.: How the National Mental Health Act works. Ment. Hyg. 33: 30; Pub. Health Rep. 64: 303.

*Lull, Clifford B. (and Hingson): Control of pain in childbirth. Ed. 3. Philadelphia, J. B. Lippineott Company, 522 pp.

M

*Magnuson, Harold J. (and Rosenau): The rate of development and degree of acquired immunity in experimental rabbit syphilis. In Recent advances in the study of venereal diseases. A symposium, Raleigh, N. C., Venereal Disease Education Institute in cooperation with the United States Public Health Service, p. 1.

* — The treatment of asymptomatic neurosyphilis in the white mouse. In Recent advances in the study of venereal diseases. A symposium, Raleigh, N. C. Venereal Disease Education Institute in cooperation with the United States Public Health Service, p. 332.

United States Public Health Service, p. 332.

Mahoney, John F.: The treatment of syphilis in the United States. The

Practitioner (London) 162: 358.

* (Wright, and Trautman): The therapeutic efficacy of aqueous penicillin in early syphilis. In Recent advances in the study of venereal diseases. A symposium, Raleigh, N. C., Venereal Disease Education Institute in cooperation with the United States Public Health Service, p. 95.

Mahoney, John F. (and Zwally): Observations on the serodiagnosis of syphilis. Bull. World Health Organization 2: 227.

See also Arnold, R. C. (Mahoney, Nicholson, and Wright).

Maier, Paul P. (and Dow): Diarrheal disease control studies. II. Conical net for collecting flies. Pub. Health Rep. 64: 604.

Mandel, Emanuel E.: Rapid test for serum protein alterations. Mil. Surgeon 104: 456.

Marks, Edna K.: See Jacobson, Leon O. (Marks and Lorenz).

Marshak, Alfred (Schaefer and Rajagopalan): Antibacterial activity of d-usnic acid and related compounds on *M. tuberculosis*. Proc. Soc. Exper. Biol. and Med. **70**: 565.

Marshall, Louise H. (and Specht): Respiratory water vapor at simulated altitude. Am. J. Physiol. 156: 299.

Marshall, Wade H.: The presence of multiple cortical foci in the primary sensory projection systems of the cat (abstract). Federation Proc. 8: 107.

———Suppressor action on primary sensory projection reactions (abstract). Federation Proc. 8: 107.

Maver, Mary E. (and Greco): Depolymerase activities of calf spleen and thymus cathepsin preparations (abstract). Federation Proc. 8: 227.

McCauley, H. Berton: Topical fluoride solutions for the control of dental caries: Considerations pertinent to their clinical application. J. Am. Dent. Assoc. 38: 430.

——— (Likins and Arnold): The relation between dental caries experience and poliomyelitis in Minneapolis school children (1946 epidemic). J. Am. Dent. Assoc. 38: 735.

McCleod, Charlotte (and Arnold): Preservation and inoculation studies on Treponema pallidum. J. Ven. Dis. Inform. 30: 104.

McClosky, W. T.: See Emmart, Emily W. (McClosky, Smith, and Lieberman). McClure, F. J.: Fluorine and other trace elements in nutrition. J. Am. Med. Assoc. 139: 711.

- See also Zipkin, I. (and McClure).

McCoy, G. W.: Modern public health measures against leprosy. Pub. Health Nursing 41: 28.

McCullough, Norman B. (and Eisele): Toxic reactions during streptomycinsulfadiazine therapy of brucellosis. J. Am. Med. Assoc. 139: 80.

——— (Eisele and Pavelcheck): Isolation of Brucella abortus from hogs. Pub. Health Rep. 64: 537.

- See also Apter, Nathaniel S. (Eisele, McCullough, and Halstead).

*McDaniel, T. W. (and Diamond): Study of 700 discharged neuropsychiatric casualties and follow-up. Dis. Nerv. System 9: 148.

McDonald, Roger K.: See Miller, John H. (McDonald and Shock).

*McGibony, J. R.: Expansion of hospital facilities. Bull. Am. Soc. Hosp. Pharm. 5: 150; Oklahoma State Hosp. J. 3: 3, May.

- See also Miller, James S. (and McGibony).

McIver, Pearl: Public health nursing responsibilities. Editorial. Pub. Health Nursing 41: 65.

Mee, Robert A.: Gonococcal infection of the median raphe. Am. J. Syph., Gonor., and Ven. Dis. 33: 182.

Meis, Armon: See Oliphant, John W. (Gordon, Meis, and Parker).

Meister, Alton: Enzymatic degradation of triacetic acid lactone determined by a spectrophotometric method. Arch. Biochem. 20: 168.

—— Metabolism of 3,5-diketohexanoic acid and its Δ-lactone by tissue homogenates. J. Biol. Chem. 178: 577.

Meister, Alton: Metabolism of 2,4- and 3,5-diketo acids (abstract). Federation Proc. 8: 228.

----- See also Price, Vincent E. (Meister and Greenstein).

Merwin, Ruth M.: See Algire, G. H. (Merwin and Legallais).

Michele, Arthur A. (and Harper): A method of spinal fusion by arthrodesis and iliac bone grafts. Mil. Surgeon 104: 90.

*_____ (and Krueger): Olecranon fractures. J. Bone and Joint Surg. 30-A: 234.

Mickelsen, Olaf: See Caster, W. O. (and Mickelsen).

Miller, James S. (and McGibony): The hospital is incomplete without a dental service unit. Mod. Hosp. 72: 74.

Miller, John H. (McDonald and Shock): Effect of bacitracin on renal function (abstract). Federation Proc. 8: 111.

Miller, Joseph M. (Favour, Wilson, and Umbarger): A plasma factor responsible for *in vitro* lysis of leukcocytes by tuberculoprotein. Proc. Soc. Exper. Biol. and Med. **70**: 738.

Moriyama, Iwao M.: See Fales, W. Thurber (and Moriyama).

Morlan, Harvey B.: See Dent, Jack E. (Morlan and Hill).

Morse, John W.: See Bowdoin, C. D. (Henderson, Davis, Remein, and Morse). Muhlethaler, Kurt: Electron micrographs of plant fibers. Biochimica et Biophysica Acta 3: 15.

See also Frey-Wyssling, A. (Muhlethaler and Wyckoff).

Mulay, A. S. (and Firminger): Induction of hepatic carcinoma in rats by 4-dimethylaminobenzene-1-azo-1-naphthalene (abstract). Federation Proc. 8: 363.

Murphy, Edwin D.: See Kaplan, Henry S. (and Murphy). Murphy, Marie K.: See Wooley, Jerald G. (and Murphy).

Musselman, A. D.: See Eagle, Harry (Fleischman and Musselman).

N

Nadel, Eli M.: Fibroids in a guinea pig (family 13) after partial castration. J. Nat. Cancer Inst. 9: 271.

——— (Taylor, Greenberg, and Josephson): Adrenal hypertrophy in chicks infected with *Plasmodium gallinaceum*. J. Nat. Malaria Soc. 8: 70.

See also Josephson, Edward S. (Taylor, Greenberg, and Nadel).

Neal, P. A.: See Spicer, Samuel S. (and Neal).

Ness, A. T.: See Endicott, K. M. (Gillman, Brecher, Ness, Clarke, and Adamik).
Newton, Walter L.: (Bennett and Figgat): Observations on the effects of various sewage treatment processes upon eggs of Taenia saginala. Am. J. Hyg. 49: 166.

Nicholson, F. P. See Arnold, R. C. (Mahoney, Nicholson, and Wright); Arnold, R. C. (and Nicholson).

0

*O'Brien, Henry R.: Doctors and nurses of other lands. Connecticut State Med. J. 12: 28.

Odoroff, Maurice E.: Vocational rehabilitation. Spastic Review 10: 11.

Vocational rehabilitation for the cerebral palsied. Nerv. Child. 8: 214. Oliphant, John W. (Gordon, Meis, and Parker): Q fever in laundry workers, presumably transmitted from contaminated clothing. Am. J. Hyg. 49: 76.

Olivier, Louis J.: Penetration of dermatitis-producing schistosome cercariae. Am. J. Hyg. 49: 134.

- Olivier, Louis J.: Schistosome dermatitis, a sensitization phenomenon. Am. J. Hyg. 49: 290.
- Olson, Byron J.: See Bell, Joseph A. (Pittman and Olson).
- Ottinger, Barbara: See Felton, Lloyd D. (Prescott, Kauffmann, and Ottinger).

P

- Paine, Robert M. (and Shock): Variability and reliability of cardiac index estimations with the high-frequency, undamped ballistrocardiograph (abstract). Federation Proc. 8: 123.
- *Pardo, Oswaldo A. (and Farmer): Acute syphilitic meningitis: A comparison of penicillin with arseno-bismuth therapy. In Recent advances in the study of venereal diseases. A symposium, Raleigh, N. C., Venereal Disease Education Institute in cooperation with the United States Public Health Service, p. 333.
- Parker, R. R. (Bell and Stoenner): Q fever. A brief survey of the problem. J. Am. Vet. Med. Assoc. 114: 124.
- See also Bell, E. J. (Parker and Stoenner); Lackman, David B. (Parker and Gerloff); Oliphant, John W. (Gordon, Meis, and Parker).
- *Parkhurst, George E.: Treatment of neurosyphilis at Hot Springs V. D. Medical Center. In Recent advances in the study of venereal diseases. A symposium, Raleigh, N. C., Venereal Disease Education Institute in cooperation with the United States Public Health Service, p. 209.
- See also Cannefax, George R. (Parkhurst and Bowman).
- Patnode, Robert A.: See Spendlove, George A. (Cummings and Patnode).
- Pelton, Walter J.: Dental caries data obtained by simplified method and correlated with fluorine data in four Wyoming cities. J. Am. Dent. Assoc. 38: 723.
- Pennell, Maryland P.: See Hubbard, John P. (Bain and Pennell).
- Perkins, Keith J.: First aid in mental health. Arizona Pub. Health News 42: 2, Jan.
- Perrin, Theodore: Histopathologic observations in a fatal case of Q fever. Arch. Path. 47: 361.
- Perrott, George St. J.: The morbidity survey in public health work. Am. J. Pub. Health 39: 741.
- Pescor, Michael J.: The Department of Public Health [Georgia] mental health program. J. Med. Assoc. Georgia 38: 119.
- A study of selective service law violators. Am. J. Psychiat. 105: 641.
- Petry, Lucile: Evolution of nurse types. Hospitals 23: 64, May.
- ---- The National Nursing Accrediting Service. Am. J. Nursing 49: 362.
- *Philip, Cornelius B. (and Kohls): Mites and scrub typhus. In Proc. 4th Internat. Cong. Trop. Med. and Malaria, Washington, D. C., U. S. Government Printing Office, vol. 2, p. 1656.
- Pittman, Margaret: See Bell, Joseph A. (Pittman and Olson); Davis, Dorland J. (and Pittman).
- Poel, William E.: Effect of anoxic anoxia on myoglobin concentration in striated muscle. Am. J. Physiol. 156: 44.
- Poindexter, Hildrus A.: Experimental animal colony in tropical West Africa. Pub. Health Rep. 64: 57.
- *Polson, A.: Quantitative partition chromatography and the composition of E. coli. Biochimica et Biophysica Acta 2: 575.
- Pond, M. Allen: Housing and health. Sanitary aspects of the dwelling. Am. J. Pub. Health 39: 454.
- Porterfield, Virginia T.: See Heppel, Leon A. (and Porterfield).
- Prescott, Benjamin: See Felton, Lloyd D. (Prescott, Kauffmann, and Ottinger).

Price, David E.: Research grants awarded by Public Health Service. Pub. Health Rep. Supp. 205 (revised), 54 pp.

Price, Eleanor V.: See Bauer, Theodore J. (and Price).

Price, Vincent E. (Meister and Greenstein): Partial separation of D- and L- peptidases and analogous dehydropeptidase (abstract). Federation Proc. 8: 238.

—— See also Fodor, Paul J. (Price and Greenstein); Greenstein, Jesse P. (and Price).

R

Rajagopalan, Srinivasa: See Marshak, Alfred (Schaefer and Rajagopalan).

Randak, Edward F.: See Boone, Bert R. (Randak, Ellinger, and Oppenheimer). Rapoport, S. (Brodsky, West, and Mackler): Urinary flow and excretion of solutes during osmotic diuresis in hydropenic man. Am. J. Physiol. 156: 433.

Reardon, Lucy V. (and Bartgis): The cultivation of the small race of *Endamoeba histolytica* with a single species of bacteria. J. Parasitol. 35: 218.

Reed, Louis S.: See Rice, Dorothy P. (and Reed).

Reid, Mary E.: Relation of age to ascorbic acid metabolism in guinea pigs (abstract). Federation Proc. 8: 392.

Rein, Charles R. (Bukantz, Kent, Cooper, Ruhe, and Coatney): Studies in human malaria. XIX. The course of the complement-fixation reaction in sporozoite-induced St. Elizabeth strain vivax malaria. Am. J. Hyg. 4: 374.

Remein, Quentin R.: See Bowdoin, C. D. (Henderson, Davis, Remein, and Morse). Renes, Lucian E.: Environmental aspects of the Industrial Hygiene Study in the Ferrous Foundries of Illinois. Preliminary report. Am. Indust. Hyg. Assoc. Quart. 10: 5.

*Reynolds, Frank W.: Penicillin in early syphilis: An analysis of the discrepancies between the results of Arnold et al. and those of the Central Statistical Unit. In Recent advances in the study of venereal diseases. A symposium, Raleigh, N. C., Venereal Disease Education Institute in cooperation with the United States Public Health Service, p. 113.

Rice, Dorothy P. (and Reed): The Nation's needs for hospitals and health centers. A summary of data from plans submitted by the States under the Hospital Survey and Construction Act. Washington, D. C., Public Health Service. Processed.

Richtmyer, Nelson K.: Functional derivatives of the antimalarial 9-(2-diamylamino-1-hydroxyethyl)-1,2,3,4-tetrahydrophenanthrene (SN 1796). J. Org. Chem. 14: 334.

- See also Stewart, Laura C. (Richtmyer and Hudson).

Riley, Vernon T. (Hesselbach, Fiala, Woods, and Burk): Application of chromatography to the separation of subcellular, enzymatically active granules. Science 109: 361.

Roberson, Thomas E.: See Benjamin, Edward A. (and Roberson).

Rodriquez, Jack (Schwemlein, Bauer, Plotke, Peters, Kendell, and Rodriquez): Fever-chemotherapy in early syphilis. J. Ven. Dis. Inform. 30: 69.

Rosenau, Barbara J.: See Magnuson, Harold J. (and Rosenau).

Rosenberg, Arthur A.: See Bossak, Hilfred N. (Rosenberg and Harris).

Ross, Mabel: Mental Health Congress looks at the family. J. Home Econ. 41:71.

—— Modifiable factors in anxiety reactions. Dis. Nerv. System 10: 146.

Pilot mental health clinic. First annual report of Prince Georges County [Maryland] Clinic. Pub. Health Rep. 64: 797.

Ruhe, David S. (Cooper, Coatney, and Josephson): Studies in human malaria. XII. The protective and therapeutic action of SN 5241 against St. Elizabeth strain vivax malaria. XIV. The ineffectiveness of colchicine (SN 12,080), SN 7266, and SN 8557 as curative agents against St. Elizabeth strain vivax malaria. XV. The therapeutic action of pamaquine (plasmochin) against St. Elizabeth vivax malaria. Am. J. Hyg. 49: 346; 361; 367.

IX. The protective and therapeutic action of SN 6911 (sontochin) against

St. Elizabeth strain vivax malaria. Am. J. Hyg. 49: 41.

Russek, Henry I. (and Zohman): Hypertensive encephalopathy and cerebral

arteriosclerosis. New York State J. Med. 49: 1411.

8

Sabine, Jean Captain: Kinetics of cholinesterase of erythrocytes in pernicious and other anemias (abstract). Federation Proc. 8: 136.

Salvin, S. B.: Cysteine and related compounds in the growth of the yeastlike phase of *Histoplasma capsulatum*. J. Infect. Dis. 84: 275.

——— Phase-determining factors in Blastomyces dermatitidis. Mycologia 41: 311.

Sanford, Katherine K.: See Earle, Wilton R. (Evans, Sanford, Likely, and Schilling).

Sargent, Lewis J.: The N-acylation of N-(4-methoxyphenl)-4-chloroanthranilic acid. J. Org. Chem. 14: 285.

Schacter, Bernard: Polyphenoloxidase activity of blood serum (abstract). Federation Proc. 8: 144.

Scheele, Leonard A.: Address of welcome. In Proc. 47th Annual Conference, Surgeon General, Public Health Service, and Chief, Children's Bureau with State and Territorial Health Officers, State Mental Health Authorities, and State Hospital Survey Authorities, Washington, D. C., Public Health Service, p. 1. Processed.

- The challenge of chronic disease. Wisconsin Med. J. 48: 437.

——— Challenge of the National Health Assembly. In Proc. Sixty-second Annual Meeting, Conference of State and Provincial Health Authorities of North America, Chicago, The Conference, p. 25.

A forward look at public health. Life and Health 64: 18, June.

—— General principles of the Federal program for mental health. Mental Hyg. 33: 25.

——— Health service for the adult population. Indust. Hyg. Newsletter 9: 7, Mar.

Medical education in programs of the Public Health Service. J. Am. Med. Assoc. 140: 5.

The medical school and national health. J. Assoc. Am. Med. Colleges 24: 87.

The past and the future of the Public Health Service. Am. J. Pub. Health 39: 293.

—— The private physician in venereal disease control. J. Nat. Med. Assoc. 41: 9.

—— Public health. In Social work yearbook, New York, Russell Sage Foundation, p. 381.

Scheele, Leonard A.: We are moving forward. A Social Hygiene Day report on progress in venereal disease control. J. Social Hyg. 35: 110.

World interests of the Public Health Service. Am. Foreign Service J. 26: 10.

Schilling, E. L.: See Earle, Wilton R. (Evans, Sanford, Likely, and Schilling).

Schmehl, Francis L.: See Smith, Nathan L. (and Schmehl).

Schneider, Walter C. (and Potter): Intracellular distribution of enzymes. IV.

The distribution of oxalacetic oxidase activity in rat liver and rat kidney fractions. J. Biol. Chem. 177: 893.

*Schumacher, Henry C.: The integration of mental hygiene concepts and practices in a public health program. Canadian J. Pub. Health 39: 351.

Schwemlein, George X.: See Rodriquez, Jack (Schwemlein, Bauer, Plotke, Peters, Kendell, and Rodriquez).

Schwartz, Louis: Dermatitis among the public from new fabrics, dyes, and finishes. Chem. and Eng. News 27: 1358.

Scott, David B. (Kaplan and Wyckoff): Replica studies of changes in tooth surfaces with age. J. Dent. Res. 28: 31.

(and Wyckoff): Metal shadowing for the optical microscopy of certain tissues. Am. J. Clin. Pathol. 19: 63.

Sebrell, William H.: Malnutrition. In Geriatric medicine. The care of the aging and the aged, edited by Edward S. Stieglitz, Philadelphia, W. B. Saunders Co. Ed. 2, p. 187.

—— See also Aykroyd, W. R. (Jolliffe, Lowry, Moore, Sebrell, Shank, Tisdall, Wilder, and Zamecnik):

*Shapiro, Herbert: The change in osmotically inactive fraction produced by cell activation. J. Gen. Physiol. 32: 43.

Shimkin, Michael B.: (and Bierman): Chymotrypsin in cancer. Proc. Soc. Exper. Biol. and Med. 71: 250.

——— (Saperstein, Goetzl, Wheeler, and Berlin): Blood histamine in leukemia and erythremia. J. Nat. Cancer Inst. 9: 379.

Shock, Nathan W.: See Cohn, Jerome E. (and Shock); Davis, James O. (and Shock); Farber, Herbert R. (Yiengst and Shock); Miller, John H. (McDonald and Shock); Paine, Robert M. (and Shock); Smith, Luther E. (and Shock); Yiengst, Marvin J. (and Shock).

Silverman, Milton: Interactions of atabrine, thiamine, and cocarboxylase. J. Biol. Chem. 178: 423.

*Simmons, Samuel W. (and Hayes): Fleas and disease. In Proc. 4th Internat. Cong. Trop. Med. and Malaria, Washington, D. C., U. S. Government Printing Office, vol. 2, p. 1678.

Skinner, Homer L.: See Iverson, Roland K. (and Skinner).

*Smith, Falconer (and Labaw): The development of a tracer method for determining the rate of diffusion of inert gas through living tissue. J. Cell. and Comp. Physiol. 32: 407.

Smith, Luther E. (and Shock): Intravenous glucose tolerance in aged males. J. Gerontol. 4: 27.

Smith, Milton I. (and Emmart): Studies in the metabolism of tubercle bacillus.

I. The production of riboflavin. J. Immunol. 61: 259.

—— (Jackson, Chang, and Longnecker): Metabolic fate of 4,4'-diaminodiphenylsulfone (DDS) in the rabbit and its isolation from urine. Proc. Soc. Exper. Biol. and Med. 71: 23.

—— See also Emmart, Emily W. (McClosky, Smith, and Lieberman); Stohlman, E. F. (Thorp and Smith).

*Smith, Nathan L. (and Schmehl): Catalytic hydrogenation and some aryl-substituted aliphatic acids. J. Org. Chem. 13: 859.

Smith, W. W.: See Blount, H. C. (and Smith).

Spear, Frances: See Habel, Karl (Endicott, Bell, and Spear).

Specht, Heinz: Physiological effects of abnormal atmospheric pressures. In Industrial hygiene and toxicology, New York, Interscience Publishers, Inc., vol. 1, p. 135.

---- See also Marshall, Louise H. (and Specht).

Spencer, R. R.: The mystery of cancer growth. Hygiea 27: 322.

—— National Cancer Institute program of postgraduate training for physicians. Pub. Health Rep. 64: 750.

Spendlove, George A. (Cummings and Patnode): Demonstration of tubercle bacilli in sputum smears. Selection of choice particles compared with various sodium hydroxide concentration techniques. Pub. Health Rep. 64: 574.

Spicer, Samuel S. (Hanna and Clark): Studies in vitro on methemoglobin reduction in dog erythrocytes. J. Biol. Chem. 177: 217.

----- (Highman and Monaco): Toxic and pathologic effects of xylidine in the fasting and non-fasting states. J. Pharm. and Exper. Therap. 95: 256.

——— (and Neal): The effect of hypoxia on the in vivo formation of methemoglobin by aniline and nitrite. J. Pharm. and Exper. Therap. 95: 438.

——— (Wooley and Clark): Spectral deviations found in hemolyzed erythrocytes of rabbits fed a purified diet (abstract). Federation Proc. 8: 148.

Spoto, Joseph F. (and Iskrant): Contact investigation of syphilis. J. Ven. Dis. Inform. 30: 140.

Spriggs, John B.: See Binford, C. H. (and Spriggs); Gledhill, Emerson Y. (Spriggs and Binford).

Stark, T. H.: See Emmart, Emily W. (Kissling and Stark).

Steele, James H. (and Tierkel): Rabies problems and control. A nation-wide program. Pub. Health Rep. 64: 785.

Steinle, John G.: See Hoge, Vane M. (and Steinle).

Stewart, H. L.: See Burgoyne, F. H. (Heston, Hartwell, and Stewart); Hare, W. V. (and Stewart).

Stewart, Laura C. (Richtmyer and Hudson): Trehalose from yeast. J. Am. Chem. Soc. 71: 2277.

Stoenner, H. G.: See Bell, E. J. (Parker and Stoenner); Parker, R. R. (Bell and Stoenner).

Stohlman, E. F. (Thorp and Smith): Pharmacologic action of chlordane (abstract). Federation Proc. 8: 334.

Strauss, R. C.: Occupational mortality rates, 1950 subject of study. Indust. Hyg. Newsletter 9: 2, Jan.

Striebich, Mary J.: See Dalton, Albert J. (Kahler, Kelly, Lloyd, and Striebich); Dalton, Albert J. (Kahler, Lloyd, and Striebich).

Sullivan, Catherine M.: No help needed [orthopedic nursing]. Am. J. Nursing 49: 90.

- Orthopedic nursing. Richmond County Med. Soc. Bull. 7: 69.

Swinney, Daniel D.: Current organizational patterns of statistical activities in State health departments. Pub. Health Rep. 64: 621.

Symeonidis, A.: Eclampsis-like condition in pregnant rats injected with progesterone (abstract). Federation Proc. 8: 371.

*——: Mammakrebserzeugung bei Mäusen durch Progesteron verabreicht während der Gravidität (Mammary tumors induced in mice injected with progesterone during pregnancy—summary). Acta Unio Internationalis Contra Cancrum 6: 163.

Szent-Györgyi, Albert: Free-energy relations and contraction of actomyosin. Biol. Bull. 96: 140.

- See also Borbiro, M. (and Szent-Györgyi).

- Tabor, Herbert: Purification of diamine oxidase (histaminase) (abstract). Federation Proc. 8: 335.
- Tauber, Henry: Isolation of a crystalline heat stable trypsin inhibitor from lima beans (abstract). Federation Proc. 8: 259.
- A new color test for tryptophan and related compounds. J. Biol. Chem. 177: 337.
- Taylor, D. Jane: See Josephson, Edward S. (Taylor, Greenberg, and Nadel); Nadel, Eli M. (Taylor, Greenberg, and Josephson).
- Terry, Luther L.: Oral aluminum penicillin, a clinical study. South. M. J. 42: 507.
- Thayer, J. D.: See Carpenter, Charles M. (Bucca, Buck, Casman, Christensen, Crowe, Drew, Hill, Lankford, Morton, Peizer, Shaw, and Thayer).
- Thorp, W. T. S.: See Stohlman, E. F. (Thorp and Smith).
- Tibbitts, Clark: Adjustment in old age—The beginning of a university experiment. Michigan Alumnus 55: 234.
- Tiboni, Emil A.: Housing and health. Appraisal of substandard housing. Am. J. Pub. Health 39: 459.
- Tierkel, Ernest S.: See Steele, James H. (and Tierkel).
- Topping, Norman H.: Research progress on the common cold. Wisconsin Med. J. 48: 418.
- ——— Rocky Mountain spotted fever. In Veterans' Administration Technical Bulletin, Washington, D. C., series 10, vol. 2 (TB 10-40), p. 1.
- Townsend, J. G.: Industrial hygiene organization and practices in the United States. *In Proc.* 9th Internat. Cong. Indust. Med., Bristol, England, John Wright and Sons, Ltd., p. 844.
- Trasko, Victoria M.: States spend more money on industrial hygiene. Indust. Hyg. Newsletter 9: 5, Jan.
- The work of State and local industrial hygiene agencies. Pub. Health Rep. 64: 471; Indust. Hyg. Newsletter 9: 7, Apr.
- Trautman, J. A. (and Giacomo): Integration of public health nursing into a Marine hospital. J. Ven. Dis. Inform. 30: 15.
- Public health nursing in a Marine hospital. Pub. Health Nursing
- See also Mahoney, John F. (Wright and Trautman).
- Tschirgi, R. D. (Gerard, Jenerick, Boyarsky, and Hearon): Metabolism of the rat spinal cord functioning in isolation (abstract). Federation Proc. 8: 166.
- Tucker, Harold A.: The effect of human cerebrospinal fluid on the dilution bioassay of penicillins G, X and K. In Recent advances in the study of venereal diseases. A symposium, Raleigh, N. C., Venereal Disease Education Institute in cooperation with the United States Public Health Service, p. 306.
- Penicillin syphilotherapy administered during pregnancy. A study of 149 pregnancies during which penicillin was given for early maternal syphilis. Am. J. Med. Sc. 217: 157.
- *—— (Farmer and Fein): Penicillin in cardiovascular syphilis. Follow-up note on thirty-two cases. In Recent advances in the study of venereal diseases. A symposium, Raleigh, N. C., Venereal Disease Education Institute in cooperation with the United States Public Health Service, p. 245.
- Tullner, William W.: See Hertz, Roy (Dhyse and Tullner); Hertz, Roy (and Tullner).

- Van Allen, Willard W.: Characteristics of commercial X-ray intensifying screens and films—IV. Pub. Health Rep. 64: 430.
- ——— Characteristics of commercial X-ray screens and films—V; VI. Pub. Health Rep. 64: 581; 706.
- Persistence of fluoroscopic screens. Pub. Health Rep. 64: 560.
- Van Slyke, C. J.: The National Heart Act. In Proc. 47th Annual Conference, Surgeon General, Public Health Service, and Chief, Children's Bureau with State and Territorial Health Officers, State Mental Health Authorities, and State Hospital Survey and Construction Authorities, Washington, D. C., Public Health Service, p. 27. Processed.
- The program of the National Heart Institute. Pharmacal Advance 180: 3.
- —— Public Health Service aid to medical education and research. J. Assoc. Am. Med. Colleges 24: 75.
- Vivian, Donald L.: A new synthesis of 2-phenazinol through the di-N-oxide. J. Am. Chem. Soc. 71: 1139.
- See also Waterman, Henry C. (and Vivian).
- Vogel, F. E.: See Davis, Dorland J. (and Vogel).
- *von Brand, Theodor: The physiology of helminth parasites in relation to disease.

 In Proc. 4th Internat. Cong. Trop. Med. and Malaria, Washington, D. C.,
 U. S. Government Printing Office, vol. 2, p. 984.
- von Oettingen, W. F.: Phenol and its derivatives: The relation between their chemical constitution and their effect on the organism. National Institutes of Health Bulletin No. 190. Washington, D. C., U. S. Government Printing Office, 408 pp.
- ——— (Donahue, Yagoda, Monaco, and Harris): Toxicity and potential dangers of cyclotrimethylene trinitramine (RDX). J. Indust. Hyg. and Toxicol. 31: 21.

. W

- Warner, Estella Ford: Heart disease control program. In Proc. 47th Annual Conference, Surgeon General, Public Health Service, and Chief, Children's Bureau with State and Territorial Health Officers, State Mental Health Authorities, and State Hospital Survey and Construction Authorities, Washington, D. C., Public Health Service, p. 34. Processed.
- Waterman, Henry C. (and Vivian): Direct ring-closure through the nitro group in certain aromatic compounds with the formation of nitrogen heterocycles; a new reaction. J. Org. Chem. 14: 289.
- Watt, James: Fly control and the acute diarrheal diseases. Bol. Ofic. san. panam. 28: 233.
- *Waugh, Richey L. (and Yancey): Carpometacarpal dislocations with particular reference to simultaneous dislocation of the bases of the fourth and fifth metacarpals. J. Bone and Joint Surg. 30-A: 397

White, Julius: See Fones, William S. (and White).

Wight, Kent (and Burk): Progesterone and testosterone reversal of diethylstilbestrol inhibition of succinoxidase of tumor mitochondrial elements (abstract). Federation Proc. 8: 265.

*Wikler, Abraham (and Frank): Effects of electroshock convulsions on chronic decorticated cats. Proc. Soc. Exper. Biol. and Med. 67: 464.

*—— (Wiesel and Maxwell): Sympathetic paralysis due to metastasis as initial sign of gastric carcinoma. Ann. Int. Med. 28: 648.

Wilcox, Aimee: See Haas, Victor H. (Wilcox and Coleman).

Wilkerson, Hugh L. C.: Diabetes as a public health problem. Nat. Negro Health News 17: 6, Jan.-June.

——— (and Ford): Diabetes control in a local health department. Am. J. Pub. Health 39: 607.

Williams, C. L.: The control of murine typhus with DDT. Mil. Surgeon 104: 163. Woods, Mark W. (DuBuy, Burk, and Hesselbach): Cytological studies on the nature of the cytoplasmic particulates in the Cloudman S91 mouse melanoma, the derived Algire S91A partially amelanotic melanoma, and the Harding-Passey mouse melanoma. J. Nat. Cancer Inst. 9: 311.

See also DuBuy, H. G. (Woods, Burk, and Lackey); Riley, Vernon T.

(Hesselbach, Fiala, Woods, and Burk).

Wooley, Jerald G. (and Murphy): Metabolic studies on T₂ Escherichia coli bacteriophage. I. The study of desoxyphyridoxine inhibition and its reversal. J. Biol. Chem 178: 869.

- See also Spicer, Samuel S. (Wooley and Clark).

Woolsey, Theodore D.: Planning the morbidity survey. Am. J. Pub. Health 39: 743.

Wright, John T.: See Bell, J. Frederick (Wright and Habel); Habel, Karl (Bell and Wright).

Wright, Robert D. See Arnold, R. C. (Mahoney, Nicholson, and Wright);
Mahoney, John F. (Wright and Trautman).

*Wright, Willard H. (and Cram): Wartime research in human schistosomiasis.

In Proc. 4th Internat. Cong. Trop. Med. and Malaria, Washington, D. C.,
U. S. Government Printing Office, vol. 2, p. 997.

Wyckoff, Ralph W. G.: Electron microscopy. Du(Zurich) 9: 48, Apr.

* The electron microscopy of macromolecular crystals. Acta Crystallo-graphica 1: 292.

Multiplication of the T3 bacteriophage against E. coli (abstract). Proc.

Soc. Exper. Biol. and Med. 71: 144.

See also Frey-Wyssling, A. (Muhlethaler and Wyckoff); Scott, David B. (Kaplan and Wyckoff); Scott, David B. (and Wyckoff).

Y

Yagoda, Herman: Photographic recording media [for radioactive radiations].
Nucleonics 4: 134.

 Radioactive measurements with nuclear emulsions, New York, John Wiley and Sons, 356 pp.

See also von Oettingen, W. F. (Donahue, Yagoda, Monaco, and Harris). Yiengst, Marvin J. (and Shock): Fasting serum vitamin A, and carotene levels and vitamin A absorption in males over 40 (abstract). Federation Proc. 8: 173.

Yiengst, Marvin J. (and Shock): See also Farber, Herbert R. (Yiengst and Shock).
 Young, LeRoy K.: Tuberculosis control in the Philippines. Annual report of the Tuberculosis Control Division, Philippine Public Health Rehabilitation Program, year ending June 30, 1948. Pub. Health Rep. 64: 123.

*Young, Martin D.: Adaptability of exotic malaria parasites to indigenous anophelines. In Proc. 4th Internat. Cong. Trop. Med. and Malaria, Washing-

ton, D. C., U. S. Government Printing Office, vol. 1, p. 672.

See also Coatney, G. Robert (Ruhe, Cooper, Josephson, and Young); Cooper, W. Clark (Ruhe, Coatney, Josephson, and Young); Cooper, W. Clark (Ruhe, Coatney, Josephson, Young, and Burgess); Ruhe, David S. (Cooper, Coatney, Josephson, and Young).

7.

Zimmer, D. J.: See Webster, Stewart H. (Liljegren and Zimmer).
Zipkin, I. (and McClure): Inhibitory effect of fluoride on tooth decalcification by citrate and lactate in vivo. J. Dent. Res. 28: 151.
Zohman, Burton L.: See Russek, Henry I. (and Zohman).
Zwally, Margaret R.: See Mahoney, J. F. (and Zwally).

0

PUBLIC LIBRARY

DETROIT

